


### Supplementary Information

**Three decades later in the northern Kruger National Park: multiple acoustic and capture surveys may underestimate the true local richness of bats based on historical collections**


**Erin R. Brinkley , Sina Monika Weier, Dan M. Parker, Peter John Taylor**

**Figures S1-S25** Sonograms of each species recorded during this study in the northern Kruger National Park, South Africa. The x-axis shows the length call; duration (ms) and the y-axis the frequency (kHz).


#### **Emballonuridae: *Taphozous mauritanus***




#### **Hipposideridae: *Hipposideros caffer***




#### **Miniopteridae: *Miniopterus natalensis***




**Molossidae:** *Chaerephon ansorgei*




*Chaerephon pumilus*











*Mops condylurus*









*Mops midas*




*Otomops martiensseni**Tadarida aegyptiaca***Rhinolophidae:** *Rhinolophus darlingi**Rhinolophus fumigatus*


*Rhinolophus simulator**Rhinolophus smithersi***Vespertilionidae:** *Eptesicus hottentotus**Myotis tricolor*

*Neoromicia capensis**Neoromicia nana**Neoromicia zuluensis*

*Nycticeinops schlieffeni**Pipistrellus hesperidus**Pipistrellus rusticus*

*Scotophilus dinganii*

## Unknown34



## Unknown75

