

## List of papers collected for the review

- Alberdi, A., Aizpurua, O., Aihartza, J. & Garin, I. (2014) Unveiling the factors shaping the distribution of widely distributed alpine vertebrates, using multi-scale ecological niche modelling of the bat *Plecotus macrobullaris*. *Frontiers in Zoology*, **11**, 77.
- Amorim, F., Carvalho, S.B., Honrado, J. & Rebelo, H. (2014) Designing optimized multi-species monitoring networks to detect range shifts driven by climate change: a case study with bats in the north of Portugal. *PLoS One*, **9**, e87291.
- Becker, N.I. & Encarnação, J.A. (2012) Cost-effectiveness of habitat-suitability maps using low-detailed data for elusive bat species. *European Journal of Wildlife Research*, **58**, 945–953.
- Bellamy, C., Scott, C. & Altringham, J. (2013) Multiscale, presence-only habitat suitability models: fine-resolution maps for eight bat species. *Journal of Applied Ecology*, **50**, 892–901.
- Bilgin, R., Kesisoglu, A. & Rebelo, H. (2012) Distribution patterns of bats in the Eastern Mediterranean Region through a climate change perspective. *Acta Chiropterologica*, **14**, 425–437.
- Dixon, M.D. (2011) Post-Pleistocene range expansion of the recently imperiled eastern little brown bat (*Myotis lucifugus lucifugus*) from a single southern refugium. *Ecology and Evolution*, **1**, 191–200.
- Dormann, C.F., Gruber, B., Winter, M. & Herrmann, D. (2010) Evolution of climate niches in European mammals? *Biology Letters*, **6**, 229–32.
- Ducci, L., Agnelli, P., Di Febbraro, M., Frate, L., Russo, D., Loy, A., Carranza, M.L., Santini, G. & Roscioni, F. (2015) Different bat guilds perceive their habitat in different ways: a multiscale landscape approach for variable selection in species distribution modelling. *Landscape Ecology*, **30**, 2047–2159.
- Elith, J., Graham, C.H., Anderson, R.P., Dudík, M., Ferrier, S., Guisan, A., Hijmans, R.J., Huettmann, F., Leathwick, J.R., Lehmann, A., Li, J., Lohmann, L.G., Loiselle, B.A., Manion, G., Moritz, C., Nakamura, M., Nakazawa, Y., Overton, J.M.M., Peterson, A.T., Phillips, S.J., Richardson, K., Scachetti-Pereira, R., Schapire, R.E., Soberon, J., Williams, S., Wisz, M.S. & Zimmermann, N.E. (2006) Novel methods improve prediction of species' distributions from occurrence data. *Ecography*, **29**, 129–151.
- Elith, J. & Leathwick, J. (2007) Predicting species distributions from museum and herbarium records using multiresponse models fitted with multivariate adaptive regression splines. *Diversity and Distributions*, **13**, 265–275.
- Escalante, T., Morrone, J.J. & Rodríguez-Tapia, G. (2013) Biogeographic regions of north american mammals based on endemism. *Biological Journal of the Linnean Society*, **110**, 485–499.
- Escalante, T., Sanchez-Cordero, V., Morrone, J.J. & Linaje, M. (2007) Deforestation affects biogeographical regionalization: a case study contrasting potential and extant distributions of Mexican terrestrial mammals. *Journal of Natural History*, **41**, 965–984.
- Escalante, T., Szumik, C. & Morrone, J.J. (2009) Areas of endemism of Mexican mammals: Reanalysis applying the optimality criterion. *Biological Journal of the Linnean Society*, **98**, 468–478.
- Di Febbraro, M., Roscioni, F., Frate, L., Carranza, M.L., De Lisio, L., De Rosa, D., Marchetti, M.

- & Loy, A. (2015) Long-term effects of traditional and conservation-oriented forest management on the distribution of vertebrates in Mediterranean forests: a hierarchical hybrid modelling approach. *Diversity and Distributions*, **21**, 1141–1154.
- Flanders, J., Wei, L., Rossiter, S.J. & Zhang, S. (2011) Identifying the effects of the Pleistocene on the greater horseshoe bat, *Rhinolophus ferrumequinum*, in East Asia using ecological niche modelling and phylogenetic analyses. *Journal of Biogeography*, **38**, 439–452.
- Fonderflick, J., Azam, C., Brochier, C., Cosson, E. & Quékenborn, D. (2015) Testing the relevance of using spatial modeling to predict foraging habitat suitability around bat maternity: A case study in Mediterranean landscape. *Biological Conservation*, **192**, 120–129.
- Garcia, R.A., Burgess, N.D., Cabeza, M., Rahbek, C. & Araújo, M.B. (2012) Exploring consensus in 21st century projections of climatically suitable areas for African vertebrates. *Global Change Biology*, **18**, 1253–1269.
- Graham, C.H., Elith, J., Hijmans, R.J., Guisan, A., Townsend Peterson, A., Loiselle, B.A., Anderson, R.P., Dudk, M., Ferrier, S., Huettmann, F., Leathwick, J., Lehmann, A., Li, J., Lohmann, L., Loiselle, B., Manion, G., Moritz, C., Nakamura, M., Nakazawa, Y., Overton, J., Phillips, S., Richardson, K., Pereira, R.S., Schapire, R., Sober??n, J., Williams, S., Wisz, M. & Zimmermann, N. (2008) The influence of spatial errors in species occurrence data used in distribution models. *Journal of Applied Ecology*, **45**, 239–247.
- Greaves, R.K., Sanderson, R. a. & Rushton, S.P. (2006) Predicting species occurrence using information-theoretic approaches and significance testing: An example of dormouse distribution in Cumbria, UK. *Biological Conservation*, **130**, 239–250.
- Hahn, M.B., Epstein, J.H., Gurley, E.S., Islam, M.S., Luby, S.P., Daszak, P. & Patz, J.A. (2014) Roosting behaviour and habitat selection of *Pteropus giganteus* reveal potential links to Nipah virus epidemiology. *Journal of Applied Ecology*, **51**, 376–387.
- Hayes, M.A., Ozenberger, K., Cryan, P.M. & Wunder, M.B. (2015) Not to put too fine a point on it — does increasing precision of geographic referencing improve species distribution models for a wide-ranging migratory bat? *Acta Chiropterologica*, **17**, 159–169.
- Herkt, K.M.B., Barnikel, G., Skidmore, A.K. & Fahr, J. (2016) A high-resolution model of bat diversity and endemism for continental Africa. *Ecological Modelling*, **320**, 9–28.
- Hernandez, P.A., Franke, I., Herzog, S.K., Pacheco, V., Paniagua, L., Quintana, H.L., Soto, A., Swenson, J.J., Tovar, C., Valqui, T.H., Vargas, J. & Young, B.E. (2008) Predicting species distributions in poorly-studied landscapes. *Biodiversity and Conservation*, **17**, 1353–1366.
- Hof, A.R., Jansson, R. & Nilsson, C. (2012) Future climate change will favour non-specialist mammals in the (Sub)Arctics. *PLoS One*, **7**, e52574.
- Hughes, A.C., Satasook, C., Bates, P.J.J., Bumrungsri, S. & Jones, G. (2012) The projected effects of climatic and vegetation changes on the distribution and diversity of Southeast Asian bats. *Global Change Biology*, **18**, 1854–1865.
- Hughes, A.C., Satasook, C., Bates, P.J.J., Soisook, P., Sritongchuay, T., Jones, G. & Bumrungsri, S. (2010) Echolocation call analysis and presence-only modelling as conservation monitoring tools for Rhinolophoid bats in Thailand. *Acta Chiropterologica*, **12**, 311–327.
- Illoldi-Rangel, P., Fuller, T., Linaje, M., Pappas, C., Sánchez-Cordero, V. & Sarkar, S. (2008) Solving the maximum representation problem to prioritize areas for the conservation of terrestrial mammals at risk in Oaxaca. *Diversity and Distributions*, **14**, 493–508.

- Illoldi-Rangel, P., Sánchez-Cordero, V. & Townsend Peterson, A. (2004) Predicting distributions of Mexican mammals using ecological niche modeling. *Journal of Mammalogy*, **85**, 658–662.
- Jaberg, C. & Guisan, A. (2001) Modelling the distribution of bats in relation to landscape structure in a temperate mountain environment. *Journal of Applied Ecology*, **38**, 1169–1181.
- Jacobs, D.S., Babiker, H., Bastian, A., Kearney, T., van Eeden, R. & Bishop, J.M. (2013) Phenotypic convergence in genetically distinct lineages of a *Rhinolophus* species complex (Mammalia, Chiroptera). *PLoS One*, **8**, e82614.
- Kuo, H.C., Chen, S.F., Fang, Y.P., Flanders, J. & Rossiter, S.J. (2014) Comparative rangewide phylogeography of four endemic Taiwanese bat species. *Molecular Ecology*, **23**, 3566–3586.
- Kusch, J. & Schmitz, A. (2013) Environmental factors affecting the differential use of foraging habitat by three sympatric species of *Pipistrellus*. *Acta Chiropterologica*, **15**, 57–67.
- Lamb, J.M., Naidoo, T., Taylor, P.J., Napier, M., Ratrimomanarivo, F.H. & Goodman, S.M. (2012) Genetically and geographically isolated lineages of a tropical bat (Chiroptera: Molossidae) show demographic stability over the late Pleistocene. *Biological Journal of the Linnean Society*, **106**, 18–40.
- Lamb, J.M., Ralph, T.M.C., Goodman, S.M., Bogdanowicz, W., Fahr, J., Gajewska, M., Bates, P.J.J., Eger, J., Benda, P. & Taylor, P.J. (2008) Phylogeography and predicted distribution of African-Arabian and Malagasy populations of giant mastiff bats, *Otomops* spp. (Chiroptera: Molossidae). *Acta Chiropterologica*, **10**, 21–40.
- Lawler, J.J., Shafer, S.L., White, D., Kareiva, P., Maurer, E.P., Blaustein, A.R. & Bartlein, P.J. (2009) Projected climate-induced faunal change in the Western Hemisphere. *Ecology*, **90**, 588–597.
- Lee, D.N., Papeş, M. & Van Den Bussche, R.A. (2012) Present and potential future distribution of common vampire bats in the Americas and the associated risk to cattle. *PLoS ONE*, **7**, e42466.
- Lisón, F. & Calvo, J.F. (2013) Ecological niche modelling of three pipistrelle bat species in semiarid Mediterranean landscapes. *Acta Oecologica*, **47**, 68–73.
- Lisón, F., Palazón, J. a. & Calvo, J.F. (2013) Effectiveness of the Natura 2000 Network for the conservation of cave-dwelling bats in a Mediterranean region. *Animal Conservation*, **16**, 528–537.
- Lundy, M.G., Buckley, D.J., Boston, E.S.M., Scott, D.D., Prodöhl, P.A., Marnell, F., Teeling, E.C. & Ian Montgomery, W. (2012) Behavioural context of multi-scale species distribution models assessed by radio-tracking. *Basic and Applied Ecology*, **13**, 188–195.
- Maiorano, L., Falcucci, A., Zimmermann, N.E., Psomas, A., Pottier, J., Baisero, D., Rondinini, C., Guisan, A. & Boitani, L. (2011) The future of terrestrial mammals in the Mediterranean basin under climate change. *Philosophical Transactions of the Royal Society of London. Series B*, **366**, 2681–92.
- Martinez, E., Peterson, a. T. & Hargrove, W.W. (2004) Ecological niches as stable distributional constraints on mammal species. *Global Ecology and Biogeography*, **13**, 305–314.
- McConville, A., Law, B.S. & Mahony, M.J. (2013) Are regional habitat models useful at a local-scale? a case study of threatened and common insectivorous bats in South-Eastern Australia. *PLoS One*, **8**.
- Milne, D.J., Fisher, A. & Pavey, C.R. (2006) Models of the habitat associations and distributions of insectivorous bats of the Top End of the Northern Territory, Australia. *Biological*

*Conservation*, **130**, 370–385.

- Mokany, K., Prasad, S. & Westcott, D.A. (2014) Loss of frugivore seed dispersal services under climate change. *Nature Communications*, **5**, 3971.
- de Moraes Weber, M. & Grelle, C.E.V. (2012) Does environmental suitability explain the relative abundance of the tailed tailless bat, *Anoura caudifer*? *Natureza e Conservacao*, **10**, 221–227.
- Moratelli, R., De Andreazzi, C.S., De Oliveira, J. a. & Cordeiro, J.L.P. (2011) Current and potential distribution of *Myotis simus* (Chiroptera, Vespertilionidae). *Mammalia*, **75**, 227–234.
- Munguía, M., Townsend Peterson, A. & Sánchez-Cordero, V. (2008) Dispersal limitation and geographical distributions of mammal species. *Journal of Biogeography*, **35**, 1879–1887.
- Newbold, T., Gilbert, F., Zalat, S., El-Gabbas, A. & Reader, T. (2009) Climate-based models of spatial patterns of species richness in Egypt's butterfly and mammal fauna. *Journal of Biogeography*, **36**, 2085–2095.
- Newbold, T., Reader, T., El-Gabbas, A., Berg, W., Shohdi, W.M., Zalat, S., El Din, S.B. & Gilbert, F. (2010) Testing the accuracy of species distribution models using species records from a new field survey. *Oikos*, **119**, 1326–1334.
- Newson, S.E., Evans, H.E. & Gillings, S. (2015) A novel citizen science approach for large-scale standardised monitoring of bat activity and distribution, evaluated in eastern England. *Biological Conservation*, **191**, 38–49.
- Pauli, B.P., Badin, H.A., Haulton, G.S., Zollner, P.A. & Carter, T.C. (2015) Landscape features associated with the roosting habitat of Indiana bats and northern long-eared bats. *Landscape Ecology*, **30**, 2015–2029.
- Peterson, A.T., Ortega-Huerta, M.A., Bartley, J., Sanchez-Cordero, V., Soberon, J., Buddenmeier, R.H., Stockwell, D.R.B., Sánchez-Cordero, V., Soberón, J., Buddemeier, R.H. & Stockwell, D.R.B. (2002) Future projections for Mexican faunas under global climate change scenarios. *Nature*, **416**, 626–629.
- Peterson, A.T., Papeş, M., Carroll, D.S., Leirs, H., Johnson, K.M. & Museum, N.H. (2007) Mammal Taxa Constituting Potential Coevolved Reservoirs of Filoviruses. *Journal of Mammalogy*, **88**, 1544–1554.
- Phillips, S.J., Dudík, M., Elith, J., Graham, C.H., Lehmann, A., Leathwick, J.R. & Ferrier, S. (2009) Sample selection bias and presence-only distribution models: implications for background and pseudo-absence data. *Ecological Applications*, **19**, 181–97.
- Rainho, A. & Palmeirim, J.M. (2013) Prioritizing conservation areas around multispecies bat colonies using spatial modeling. *Animal Conservation*, **16**, 438–448.
- Ratrimomanarivo, F.H., Goodman, S.M., Stanley, W.T., Naidoo, T., Taylor, P.J. & Lamb, J.M. (2009) Geographic and Phylogeographic Variation in *Chaerephon leucogaster* (Chiroptera: Molossidae) of Madagascar and the Western Indian Ocean Islands of Mayotte and Pemba. *Acta Chiropterologica*, **11**, 25–52.
- Razgour, O. (2015) Beyond species distribution modelling: A landscape genetics approach to investigating range shifts under future climate change. *Ecological Informatics*, **30**, 250–256.
- Razgour, O., Hanmer, J. & Jones, G. (2011) Using multi-scale modelling to predict habitat suitability for species of conservation concern: The grey long-eared bat as a case study. *Biological Conservation*, **144**, 2922–2930.

- Razgour, O., Juste, J., Ibáñez, C., Kiefer, A., Rebelo, H., Puechmaille, S.J., Arlettaz, R., Burke, T., Dawson, D.A., Beaumont, M. & Jones, G. (2013) The shaping of genetic variation in edge-of-range populations under past and future climate change. *Ecology Letters*, **16**, 1258–1266.
- Razgour, O., Rebelo, H., Puechmaille, S.J., Juste, J., Ibáñez, C., Kiefer, A., Burke, T., Dawson, D.A. & Jones, G. (2014) Scale-dependent effects of landscape variables on gene flow and population structure in bats. *Diversity and Distributions*, **20**, 1173–1185.
- Razgour, O., Salicini, I., Ibáñez, C., Randi, E. & Juste, J. (2015) Unravelling the evolutionary history and future prospects of endemic species restricted to former glacial refugia. *Molecular Ecology*, **24**, 5267–5283.
- Rebelo, H., Froufe, E., Brito, J.C.J.C., Russo, D., Cistrone, L., Ferrand, N. & Jones, G. (2012a) Postglacial colonization of Europe by the barbastelle bat: agreement between molecular data and past predictive modelling. *Molecular Ecology*, **21**, 2761–2774.
- Rebelo, H., Froufe, E., Ferrand, N. & Jones, G. (2012b) Integrating molecular ecology and predictive modelling: Implications for the conservation of the barbastelle bat (*Barbastella barbastellus*) in Portugal. *European Journal of Wildlife Research*, **58**, 721–732.
- Rebelo, H. & Jones, G. (2010) Ground validation of presence-only modelling with rare species: a case study on barbastelles *Barbastella barbastellus* (Chiroptera: Vespertilionidae). *Journal of Applied Ecology*, **47**, 410–420.
- Rebelo, H., Tarroso, P. & Jones, G. (2010) Predicted impact of climate change on European bats in relation to their biogeographic patterns. *Global Change Biology*, **16**, 561–576.
- da Rocha, P.A., Ferrari, S.F., Feijó, A. & Gouveia, S.F. (2015) Zoogeography of South American forest-dwelling bats: Disjunct distributions or sampling deficiencies? *PLoS One*, **10**, 1–10.
- Roscioni, F., Rebelo, H., Russo, D., Carranza, M.L., Di Febbraro, M. & Loy, A. (2014) A modelling approach to infer the effects of wind farms on landscape connectivity for bats. *Landscape Ecology*, **29**, 891–903.
- Roscioni, F., Russo, D., Di Febbraro, M., Frate, L., Carranza, M.L. & Loy, A. (2013) Regional-scale modelling of the cumulative impact of wind farms on bats. *Biodiversity and Conservation*, **22**, 1821–1835.
- Russo, D., Di Febbraro, M., Cistrone, L., Jones, G., Smeraldo, S., Garonna, A.P. & Bosso, L. (2015) Protecting one, protecting both? Scale-dependent ecological differences in two species using dead trees, the rosalia longicorn beetle and the barbastelle bat. *Journal of Zoology*, **297**, 165–175.
- Rutishauser, M.D., Bontadina, F., Braunisch, V., Ashrafi, S. & Arlettaz, R. (2012) The challenge posed by newly discovered cryptic species: disentangling the environmental niches of long-eared bats. *Diversity and Distributions*, **18**, 1107–1119.
- Sanchez-Cordero, V., Illoldi-Rangel, P., Linaje, M., Sarkar, S. & Peterson, A.T. (2005) Deforestation and extant distributions of Mexican endemic mammals. *Biological Conservation*, **126**, 465–473.
- Santos, H., Juste, J., Ibáñez, C., Palmeirim, J.M., Godinho, R., Amorim, F., Alves, P., Costa, H., de Paz, O., Pérez-Suarez, G., Martínez-Alos, S., Jones, G. & Rebelo, H. (2014) Influences of ecology and biogeography on shaping the distributions of cryptic species: three bat tales in Iberia. *Biological Journal of the Linnean Society*, **112**, 150–162.
- Santos, H., Rodrigues, L., Jones, G. & Rebelo, H. (2013) Using species distribution modelling to

predict bat fatality risk at wind farms. *Biological Conservation*, **157**, 178–186.

Sattler, T., Bontadina, F., Hirzel, A.H. & Arlettaz, R. (2007) Ecological niche modelling of two cryptic bat species calls for a reassessment of their conservation status. *Journal of Applied Ecology*, **44**, 1188–1199.

Schoeman, M.C., Cotterill, F.P.D. (Woody), Taylor, P.J. & Monadjem, A. (2013) Using potential distributions to explore environmental correlates of bat species richness in southern Africa : Effects of model selection and taxonomy. *Current Zoology*, **59**, 279–293.

Schoeman, M.C., Goodman, S.M., Ramasindrazana, B. & Koubínová, D. (2015) Species interactions during diversification and community assembly in Malagasy *Miniopterus* bats. *Evolutionary Ecology*, **29**, 17–47.

Soto-Centeno, J.A. & Steadman, D.W. (2015) Fossils reject climate change as the cause of extinction of Caribbean bats. *Scientific Reports*, **5**, 7971.

Stoffberg, S., Schoeman, M.C. & Matthee, C.A. (2012) Correlated genetic and ecological diversification in a widespread southern african horseshoe bat. *PLoS One*, **7**, e31946.

Struebig, M.J., Wilting, A., Gaveau, D.L.A., Meijaard, E., Smith, R.J., Fischer, M., Metcalfe, K., Kramer-Schadt, S., Abdullah, T., Abram, N., Alfred, R., Ancrenaz, M., Augeri, D.M., Belant, J.L., Bernard, H., Bezuijen, M., Boonman, A., Boonratana, R., Boorsma, T., Breitenmoser-Würsten, C., Brodie, J., Cheyne, S.M., Devens, C., Duckworth, J.W., Duplaix, N., Eaton, J., Francis, C., Fredriksson, G., Giordano, A.J., Gonner, C., Hall, J., Harrison, M.E., Hearn, A.J., Heckmann, I., Heydon, M., Hofer, H., Hon, J., Husson, S., Khan, F.A.A., Kingston, T., Krebs, D., Lammertink, M., Lane, D., Lasmana, F., Liat, L.B., Lim, N.T.L., Lindenborn, J., Loken, B., Macdonald, D.W., Marshall, A.J., Maryanto, I., Mathai, J., McShea, W.J., Mohamed, A., Nakabayashi, M., Nakashima, Y., Niedballa, J., Noerfahmy, S., Persey, S., Peter, A., Pieterse, S., Pilgrim, J.D., Pollard, E., Purnama, S., Rafiastanto, A., Reinfelder, V., Reusch, C., Robson, C., Ross, J., Rustam, R., Sadikin, L., Samejima, H., Santosa, E., Sapari, I., Sasaki, H., Scharf, A.K., Semiadi, G., Shepherd, C.R., Sykes, R., Van Berkel, T., Wells, K., Wielstra, B. & Wong, A. (2015) Targeted conservation to safeguard a biodiversity hotspot from climate and land-cover change. *Current Biology*, **25**, 372–378.

Teixeira, T.S.M., Weber, M.M., Dias, D., Lorini, M.L., Esbérard, C.E.L., Novaes, R.L.M., Cerqueira, R. & Vale, M.M. (2014) Combining environmental suitability and habitat connectivity to map rare or Data Deficient species in the Tropics. *Journal for Nature Conservation*, **22**, 384–390.

Thomassen, H. a., Fuller, T., Buermann, W., Milá, B., Kieswetter, C.M., Jarrín-V., P., Cameron, S.E., Mason, E., Schweizer, R., Schlunegger, J., Chan, J., Wang, O., Peralvo, M., Schneider, C.J., Graham, C.H., Pollinger, J.P., Saatchi, S., Wayne, R.K. & Smith, T.B. (2011) Mapping evolutionary process: A multi-taxa approach to conservation prioritization. *Evolutionary Applications*, **4**, 397–413.

Thuiller, W., Araújo, M.B. & Lavorel, S. (2004) Do we need land-cover data to model species distributions in Europe? *Journal of Biogeography*, **31**, 353–361.

Thuiller, W., Lavergne, S., Roquet, C., Boulangéat, I., Lafourcade, B. & Araujo, M.B. (2011) Consequences of climate change on the tree of life in Europe. *Nature*, **470**, 531–534.

van Toor, M.L., Jaberg, C. & Safi, K. (2011) Integrating sex-specific habitat use for conservation using habitat suitability models. *Animal Conservation*, **14**, 512–520.

Tsoar, A., Allouche, O., Steinitz, O., Rotem, D. & Kadmon, R. (2007) A comparative evaluation of

presence-only methods for modelling species distribution. *Diversity and Distributions*, **13**, 397–405.

Warren, D.L., Wright, A.N., Seifert, S.N. & Shaffer, H.B. (2014) Incorporating model complexity and spatial sampling bias into ecological niche models of climate change risks faced by 90 California vertebrate species of concern. *Diversity and Distributions*, **20**, 334–343.

Wordley, C.F.R., Sankaran, M., Mudappa, D. & Altringham, J.D. (2015) Landscape scale habitat suitability modelling of bats in the Western Ghats of India: bats like something in their tea. *Biological Conservation*, **191**, 529–536.