



Available online at:

<http://www.italian-journal-of-mammalogy.it>

doi:10.4404/hystrix-00258-2020

## Research Article

## Size shifts in late Middle Pleistocene to Early Holocene *Sus scrofa* (Suidae, Mammalia) from Apulia (southern Italy): ecomorphological adaptations?

Alessio IANNUCCI<sup>1,\*</sup>, Raffaele SARDELLA<sup>1,2</sup>, Flavia STRANI<sup>1,2,3</sup>, Beniamino MECOZZI<sup>1</sup><sup>1</sup>Dipartimento di Scienze della Terra, Sapienza Università di Roma, PaleoFactory, p.le Aldo Moro 5, I-00185, Roma, Italy<sup>2</sup>Istituto Italiano di Paleontologia Umana, via U. Aldrovandi 18, I-00197 Roma, Italy<sup>3</sup>Departamento de Ciencias de la Tierra, Área de Paleontología, Universidad de Zaragoza, Pedro Cerbuna 12, S-50009 Zaragoza, Spain**Keywords:**

Aurelian  
Bergmann's rule  
body mass  
Euro-Mediterranean  
large mammals  
Quaternary

**Article history:**

Received: 15 November 2019

Accepted: 23 January 2020

**Acknowledgements**

The authors are thankful for granting access to the material, the kindness and availability of: Donato Coppola (MPCCSM); Maria Carmela Del Re (MPUN); Luciano Bruni and Stefano Grimaldi (IsIPU); Linda Riti and Michele Lustrino (MUST); Riccardo Castiglia (MACUS); Rosanna Largiogna (SMCC); Elisabetta Cioppi and Luca Bellucci (IGF); Francesca Alhaque (MUCIV); Addolorata Mazzotta and Carlo Viva (ITCG); Marco Pavia (MGP). We wish to thank J. Conti, D.A. Iurino, and A. Profico for their useful suggestions and help. We thank also H. Baills and P. Magniez for their support and suggestions. We are grateful for the insightful review of the associate editor P. Raia, M. Cherin, and an anonymous reviewer. This research was funded by Sapienza University of Rome "Grandi Scavi" 2016–2017–2018–2019 grants (principal investigator RS).

**Abstract**

The extant wild boar *Sus scrofa* has one of the largest geographical range of all mammals, and from its appearance in the late Early Pleistocene (Epivillafranchian) it is also widely represented in the European fossil record. Early forms of the species were larger than Late Pleistocene ones, but neither the chronology nor the causes of the size reduction have been thoroughly investigated. Here, we considered for the first time a large number of fossils from several late Middle Pleistocene to Early Holocene sites of the Apulian region (Italian Peninsula). In contrast to the supposed existence of a progressive trend towards small dimensions, morphometric comparisons and body mass estimates allow us to recognize several size oscillations during the late Middle Pleistocene-Early Holocene, with large forms occurring during interglacial stages and smaller ones during glacial stages. This suggests that fossil Apulian wild boar did not conform to Bergmann's rule, that predicts larger size in colder climates due to the selective pressure towards lower surface area to volume ratio. Climate recrudescence may have played an indirect role in reducing the availability of trophic resources and hence promoting the observed pattern.

**Introduction**


The extant wild boar (*Sus scrofa*) is one of the widest-ranging mammals in the world, occurring natively throughout the Palaearctic, northern Africa, the Middle-East and part of South-East Asia (Groves, 1981). Accordingly, it displays a great morphological diversity, with a significant amount of variation explained by geographical clines (Albarella et al., 2009; Groves, 2007). The species originated in Asia during the Early Pleistocene and spread in Europe during the late Early Pleistocene (Epivillafranchian) (Iannucci et al., 2020; Cherin et al., 2018; Frantz et al., 2016; Bellucci et al., 2015; Martínez-Navarro et al., 2015). It is first recorded in Europe from the ~1.1 Ma site of Untermaßfeld (Germany) (Guérin and Faure, 1997), and in Italy from the slightly younger Slivia at ~0.9–0.85 Ma (Petronio et al., 2011; Ambrosetti et al., 1979). Early forms of wild boars are often ascribed to a distinct subspecies, *S. scrofa priscus* (Goldfuss, 1823), characterized by low molar complexity and large size (Tsoukala and Guérin, 2016; Petronio et al., 2011; Guérin and Faure, 1997; Faure and Guérin, 1983; Hünermann, 1975, 1969). The validity of this taxon is often questioned, but there is a general consensus in recognizing the large dimensions of early wild boars (Made et al., 2017; Lister et al., 2010). Yet, the tempo and mode of the size reduction of this species have not been carefully investigated so far. The evolution of the wild boar is often depicted as a pro-

gressive trend towards smaller dimensions, and successively smaller chronosubspecies are sometimes identified (Hünermann, 1969), eventually leading to the modern *S. s. scrofa* since the Late Pleistocene (Faure and Guérin, 1983). However, this scenario is in contradiction with the high adaptive plasticity exhibited by the species (Ballari and Barrios-Garcia, 2014), which displays substantial variation in size and morphology throughout its geographic range (Albarella et al., 2009; Genov, 1999).

In Italy — an especially suitable territory for biochronological and paleoenvironmental studies of the Mediterranean area (Sardella et al., 2018a; Azzaroli, 1977) — Suidae fossils of large size are known from numerous Middle Pleistocene localities, such as Bristie 1 (Lugli and Sala, 2000), Cerè (Fabiani, 1919), La Polledrara di Cecanibbio (Anzidei et al., 2012), Malagrotta (Caloi and Palombo, 1979) and Torre del Pagliaccetto (Caloi and Palombo, 1978). Other scanty and/or poorly described remains come from several key localities of the Italian large mammal biochronological scale, for instance Ponte Galeria (Petronio and Sardella, 1999), Isernia La Pineta (Sala, 1996), Fontana Ranuccio (Strani et al., 2018) and Visogliano (Abbazzi et al., 2000). However, large specimens have also been recovered from Late Pleistocene sites of Italy (e.g. Minieri et al., 1995; Mirigliano, 1942; Del Campana, 1909), and more generally in Europe (e.g. Lister et al., 2010; Moigne et al., 2006; Faure and Guérin, 1983), thus challenging the hypothesis of a gradual size reduction.

\*Corresponding author


Email address: [alessio.iannucci@uniroma1.it](mailto:alessio.iannucci@uniroma1.it) (Alessio IANNUCCI)



**Figure 1** – Location of Apulia (grey area) along the Italian Peninsula. The Gargano, Murge and Salento areas are indicated. 1 – Grotta Paglicci; Murge: 2 – Grotta delle Mura; 3 – Grotte della Jena; 4 – Grotta Laceduzza; 5 – SMA-esterno; 6 – Avetranà; 7 – Grotta Mario Bernardini; Grotta Uluzzo C; 8 – Melpignano; San Sidero; 9 – Grotta delle Striare; Grotta Romanelli; Grotta Zinzulusa; 10 – Fondo Focene; II – Grotta delle Prazziche; 12 – Grotta dei Giganti; Grotta delle Tre Porte.

The existence of ecological and dimensional clines over the extant *S. scrofa* geographical range, urges to investigate its evolution excluding the confounding factor of topographical distance, which is known to affect the morphology of the species (Iannucci et al., 2020; Albarella et al., 2009; Groves, 2007, 1981; Guérin and Faure, 1997).


In this work, a comprehensive account of wild boar material from the Apulian region, in many cases reported but not described, is presented (e.g. Grotta del Cavallo, Sarti et al., 1998; Grotta Uluzzo C, Borzatti von Löwenstern and Magaldi, 1969; Grotta dei Giganti, Blanc,



**Figure 2** – Fossil specimens from Apulian sites: a – MPND683, right hemimandible from Melpignano; b – A8-II-205, left maxillary from Avetranà; c – ANDI694, left hemimandible from Avetranà; d – ANDI695, right hemimandible from Avetranà; e – ISIPU-GZ482, right hemimandible from Grotta Zinzulusa; f – MUCIV-GR3499, left hemimandible from Grotta Romanelli. Fossils are figured in labial (1), lingual (2), and occlusal (3) view. Scale bars 3 cm.

**Table 1** – Fossil material of *Sus scrofa* from the late Middle Pleistocene-Early Holocene of the Apulian Peninsula: 1 – San Sidero (1a - Cava L, 1b - fessura I); 2 – Melpignano (2a - Cava Bianco, 2b - Cava Nuzzo, 2c - Collezione Mirigliano); 3 – Grotta delle Striare; 4 – Grotta delle Tre Porte (Antro del Bambino, level F); 5 – Avetranà (5a - beds 2-7, 5b - bed 8); 6 – Grotta Zinzulusa (6a - levels B3-4, 6b - level A8); 7 – Grotta Uluzzo C (7a - level G, 7b - level F, 7c - level E); 8 – Grotta dei Giganti (levels 3f-3c); 9 – Grotta Laceduzza; 10 – Grotta Zinzulusa (10a - level B6, 10b - levels B5-3); II – SMA-esterno (IIa - level 8, IIb - levels 6A-4C); 12 – Fondo Focene; 13 – Grotta Romanelli (levels E-A); 14 – Grotta della Jena; 15 – Grotta delle Prazziche.

Anatomical portion	1a	1b	2a	2b	2c	3	4	5a	5b	6a	6b	7a	7b	7c	8	9	10a	10b	11a	11b	12	13	14	15		
Cranium									1										1			10				
Maxillary									1										1			2		8		
Upper tooth	1		1	1				2	3	8		1		1			2	2	1	1	1	6		7		
Hemimandible			5	1	1					5		1		1					2		1	5		9		
Lower tooth	1	2	2	3	1	1	3	2	8			1	2	1	2	4	3	4	1	1	3	10	1	16		
Scapula																						2				
Vertebra			15																			2				
Rib				1																						
Pelvis	2		5																2			1		1		
Humerus			1							2		1								2			1			
Radius			1		1																	1		1		
Ulna			2									1												1		
Metacarpal			1	1					1														4			
Carpal												5								1						
Femur			1									2								1			1			
Tibia			3	1								1								1						
Fibula																						2				
Metatarsal			2												1	1				1			5			
Metapodial										1	4									1			3			
Calcaneus					1						3															
Talus			3								3															
Tarsal												2												1		
I Phalanx	1							2	2										3			3				
II Phalanx	1									5	1	1								1			2			
III Phalanx											3												3			
Total		6	43	8	3	5	8	57	1	4	2	5	1	2	4	6	6	14	3	6	64	1	40			




**Figure 3** – The variation in size in *Sus scrofa*, as indicated by the length of the upper first (a), second (b), and third (c) molar. Age in millions of years (vertical axis) and oxygen isotope curves (horizontal axis; after Shackleton, 1995) on the left, followed by the fossiliferous sites and their suggested ages. Symbol: circle – Middle Pleistocene; square – Late Pleistocene-Holocene.

vestigated by performing tooth and postcranial measurements and body mass estimates.

#### Confounding factors of variations

Body size may be influenced by several factors which are not related to ecomorphological adaptations and may introduce a bias in our analysis. Here they are briefly discussed.

Although wild boars are sexually dimorphic animals with males larger than females, the sex determination of isolated remains is possible only for a few anatomical elements, namely canines and related cranial regions (e.g. mandibular symphysis, canine flanges). However, sexual size dimorphism is very low in cheek tooth measurements in *S. scrofa* (2–7%, Kuşatman, 1991; 3%, Made, 1991), and similar values occur in



**Figure 4** – The variation in size in *Sus scrofa*, as indicated by the length of the lower first (a), second (b), and third (c) molar. Age in millions of years (vertical axis) and oxygen isotope curves (horizontal axis; after Shackleton, 1995) on the left, followed by the fossiliferous sites and their suggested ages. Symbol: circle – Middle Pleistocene; square – Late Pleistocene-Holocene.

other extant suid species (2–5%, Made, 1991), whereas differences in the postcranial skeleton of *S. scrofa* are more variable and pronounced (1–16%, Kuşatman, 1991). A divergence between sexes arises during growth, with females gaining relatively less weight than males from the age of ~18–24 months, mainly as a result of the pregnancy effort (Gallo-Orsì et al., 1995).


Human populations may have played a role in the prey selection, for instance preferably hunting wild boar juveniles, as reported for the *terre brune* of Grotta Romanelli (Tagliacozzo, 2003). However, a high percentage of young individuals also occurs in sites not associated with evidence of human presence (e.g. Melpignano, 16.3%), and it more likely reflects a natural fallout of the r-strategy adopted by the species (peak in juveniles mortality, Martínez-Navarro et al., 2015; Gaillard et al., 1989).

Similarly, hunting and predatory pressure have been demonstrated to be secondary factors in limiting the population density (strictly related to body mass) of extant wild boars, in comparison with temperature and trophic resource availability (Fraendorf et al., 2016; Nores et al., 2008; Okarma et al., 1995).

Therefore, even though it is not possible to completely exclude an influence of the aforementioned factors on body size shifts, the selective pressures of the physical environment have surely played a much more important role.

#### The Apulian region

The Italian territory represents a crossroads in the Mediterranean area, playing a pivotal role in our comprehension of palaeoenvironmental dynamics during the Quaternary (Sardella et al., 2018a). Its vast latitudinal extension causes a strong climatic gradient, which in turn produces regional differences in palaeontological faunal assemblages (Petronio et al., 2007; Sala and Masini, 2007; Bedetti et al., 2001; Gliozzi et al.,



**Figure 5** – (a) Boxplot of M₃L from the late Middle Pleistocene-Early Holocene sites grouped for chronology (MIS) (see Tab. 2); (b) Table of p-values for the linear model.

**Table 2** – Chronology of the fossil material of *Sus scrofa* from the late Middle Pleistocene-Early Holocene of the Apulian Peninsula.

Site	Reference	MIS	Date
San Sidero	Mecozzi et al., 2019a; Sala et al., 1992; This work	?	Biochronology
Melpignano	Mecozzi et al., 2019a; Sala et al., 1992; This work	?	Biochronology
Grotta delle Striare	Maviglia, 1955; Cigna and Orlando, 1954; This work	?	Biochronology
Grotta delle Tre Porte	Blanc, 1958; This work	?	Biochronology
Avetrana beds 2–7	Petronio et al., 2008; Sardella et al., 2005	5	Biochronology
Grotta dei Giganti	Aureli and Ronchitelli, 2018; Blanc, 1958	4	Biochronology and Chrono-cultural
Grotta Zinzulusa - B6	Blanc, 1962; Cardini, 1962	4	Biochronology and Chrono-cultural
Grotta Mario Bernardini	Aureli and Ronchitelli, 2018; This work	4	Stratigraphy and Chrono-cultural
Grotta Uluzzo C	Aureli and Ronchitelli, 2018; This work	4	Stratigraphy and Chrono-cultural
Grotta Laceduzza	Mecozzi et al., 2019b	4	Biochronology and Chrono-cultural
Avetrana bed 8	Salari et al., 2019; Mecozzi and Bartolini Lucenti, 2018	3	Biochronology
Fondo Focene	Cancellieri, 2017	2	Biochronology and Chrono-cultural
Grotta Zinzulusa - B5–3	Blanc, 1962; Cardini, 1962	2	Biochronology and Chrono-cultural
Grotta Paglicci	Boscato, 1994	2	Radiometric data
SMA-esterno	Coppola et al., 2017; Baills, 2015	2	Radiometric data
Grotta Romanelli	Calcagnile et al., 2019; Sardella et al., 2018b	1	Radiometric data
Grotta della Jena	Anelli, 1956; This work	1	Biochronology and Stratigraphy
Grotta delle Mura	Bon and Boscato, 1993	1	Radiometric data
Grotta delle Prazziche	Borzatti von Löwenstern, 1969, 1966	1	Stratigraphy and Chrono-cultural

1997; Sala et al., 1992; Masini et al., 1990). In this context, the Apulian region (southern Italy), spanning more than 350 km from the Northwest to Southeast extremities, and with a least extension of 30 km along the Southwest-Northeast axis, represents a noteworthy case, a peninsula in the peninsula (Fig. 1).

Moreover, this territory is rich in mammal remains often associated to Mousterian technoculture (Middle Palaeolithic) and Neanderthal fossils (e.g. Fondo Cattìe, Corridi, 1989; Grotta delle Tre Porte, Blanc, 1962; see also Romandini et al., 2019). The human occupation of the region is also documented through the whole Upper Palaeolithic, with bone and lithic industry, mobiliary and rock art, and fossils of *Homo sapiens* (e.g. Grotta Romanelli, Sardella et al., 2019; Grotta Paglicci, Berto et al., 2017).

Since the end of 1800s Apulia represents a key region to study the mammal fauna evolution and the climatic changes that took place during the Quaternary in the Mediterranean area (e.g. Blanc, 1920; Botti, 1890, 1874a,b). Indeed, several Apulian fossiliferous localities are a reference point for the Italian large mammal biochronological scale, especially for the late Middle Pleistocene-Late Pleistocene transition (Petronio et al., 2007; Bedetti et al., 2001; Sala et al., 1992; among others). However, due to their complex depositional settings (inland karst systems and coastal caves), only a few Apulian sites have been recently radiometrically dated (e.g. Grotta del Cavallo, Zanchetta et al., 2018; Grotta Paglicci, Berto et al., 2017; Boscato, 1994). The earliest Apulian sites considered herein have long been chronologically attributed to the early Late Pleistocene (Sardella et al., 2005; Bologna et al., 1994; Sala et al., 1992), even though the ongoing revision of the stratigraphic and fossil data does not exclude an even older age (see Sardella et al., 2018b for discussion). For instance, the large mammal assemblage from the *terre rosse* of Melpignano was chronologically referred to the early Late Pleistocene (MIS 5). Nevertheless, the revision of the old fossil collections and the description of the new ones led to extend the biochronological time-span of this locality to the late Middle Pleistocene (Mecozzi et al., 2019a). Another Apulian site of debated chronology is Avertrana. The entire faunal assemblage has been firstly attributed to the early Late Pleistocene (MIS 5) (Pandolfi et al., 2013, 2011; Petronio et al., 2008). Later, the revision of the material from the uppermost fossiliferous layer of the succession (bed 8) led to refer this level to MIS 3 (Salari et al., 2019; Mecozzi and Bartolini Lucenti, 2018). However, there is neither a radiometric age for the deposit, nor a certain stratigraphic provenience for all the remains. If the revision of the material from bed 8 proposed a younger age for the upper part of the succession, on the other hand, the study in progress on the faunal

remains from the lower part of the deposit would suggest an older age, probably referable to the late Middle Pleistocene.

Finally, the Apulian region has also been regarded as a refuge area for several mammals adapted to warm conditions, especially during MIS 4, even though the debate on this matter still goes on (Sardella et al., 2018b; Mecozzi et al., 2018; Pandolfi and Petronio, 2015; Braun and Palombo, 2012; among others).

## Materials and methods

The sample of Suidae analysed herein is housed in several Italian institutions and museums (Tab. 1 and online resource Supplemental S1). We built our dataset by considering the available studies on specimens from the late Middle Pleistocene to the Early Holocene of the Apulian region (Tab. 2 and online resource Supplemental S1). We compared the fossil remains with a sample of extant Italian wild boars stored in the Comparative Anatomy Museum “Battista Grassi”, Sapienza University of Rome (MACUS), as well as literature data from Kuşatman (1991). Measurements of the specimens considered in this study are provided in Tab. 3 (upper teeth), Tab. 4 (lower teeth), and Tab. 5 (postcranial). Craniodental and postcranial measurements were taken to the nearest 0.1 mm with a digital caliper, as reported by Driesch (1976).

We also performed body mass estimates (Tab. 5 and 6) of the fossil and extant samples following regression equations based on the length of each molar (Damuth, 1990), on the antero-posterior distal diameter (Bd) of humerus and radius (Scott, 1990), and on the greatest length (GL) of the talus (Tsubamoto, 2014). Slope and intercept values are given in Tab. 7. Since in our sample males and females are treated together, the estimated body masses should be regarded as approximate mean values between sexes.

We further assessed the degree of intergroup affinity of the Apulian wild boar by conducting statistical analysis on the length of the lower third molar ( $M_3L$ ), which is the best represented tooth in our sample and the least affected by wear, thus the most reliable for comparisons. The dataset has been grouped according to the chronology (MIS) in order to assess the correlation between climate and size. We acknowledge that during an isotope stage several climatic oscillations may occur, but due to the lack of radiometric data the representativeness of MIS groups is usually considered an acceptable assumption, common to several studies focused on intraspecific size comparisons in the Pleistocene (e.g. Marciszak et al., 2019; Mazza and Bertini, 2013; Asperen, 2010). Moreover, this resolution is at a finer scale than what adopted in previous works on *S. scrofa* (Tsoukala and Guérin, 2016; Guérin and Faure, 1997; Faure and Guérin, 1983). MIS 1 is technically an interglacial stage, but the sites considered herein actually date to the early

Site	Reference	MIS	P <sup>3</sup> L	P <sup>3</sup> W	P <sup>4</sup> L	P <sup>4</sup> W	M <sup>1</sup> L	M <sup>1</sup> W	M <sup>2</sup> L	M <sup>2</sup> W	M <sup>3</sup> L	M <sup>3</sup> W
Melpignano	Bolognesi, 1992	?9-8									38.5	19.0
Grotta delle Tre Porte	This work	?7-5									37.4	21.9
Avetrana 2-7	This work	5									37.8	20.5
Grotta Zinzulusa - B6											(min=37.1- max=38.6)	(min=19.0- max=22.0)
Avetrana 8	This work	4										
Fondo Focone	This work	3	13.0 (min=12.2- max=13.9)	13.4 (min=12.7- max=14.1)	15.6	14.6	24.8 (min=24.0- max=25.7)	18.1 (min=16.7- max=19.6)	37.0 (min=34.5- max=39.1)	21.0 (min=19.0- max=22.0)		
SMA-esterno - Gravettian	This work	2										
SMA-esterno - Epigravettian	This work	2										
Grotta Romanelli	This work	2										
Grotta delle Mura	This work	1	14.9	10.5	13.3	13.6						
Grotta delle Prazziche	This work	1										
Extant <i>Sus scrofa</i>	Kuşatman, 1991; This work	14.5	10.9	14.2	14.4	14.5	17.7 (min=17.1- max=18.6)	14.2 (min=13.8- max=15.1)	21.5	16.4	33.9 (min=32.5- max=34.2)	18.5 (min=18.7- max=19.0)

**Table 3** – Measurements (mm) of the upper teeth of *Sus scrofa* from the late Middle Pleistocene-Early Holocene sites of Apulia and the extant Italian wild boar.

part of the Holocene, before the onset of the present-day climate regime (Giustini et al., 2018; Martrat et al., 2014; Sadoni and Narcisi, 2001). Therefore, we refer to interglacial periods (MIS 5 and MIS 3) and Early Holocene (MIS 1) separately. We evaluated differences in M3L in the late Middle Pleistocene, Late Pleistocene, Early Holocene, and extant Italian samples, using linear model with corner point parameterization testing the null hypothesis of no-differences between the mean of the reference sample (MIS 1) and the other populations ( $p>0.05$ ). Sites of uncertain age are not included in the analysis. The statistical analysis was performed using the R software (R Core Team, 2019).

## Institutional abbreviations

IGF: Museum of Natural History of the University of Florence, section of Geology and Paleontology. IIPP: Italian Institute of Prehistory and Protohistory. IsIPU: Italian Institute of Human Paleontology. ITCGC: Commercial Technical Institute “Galilei-Costa”. MACUS: Comparative Anatomy Museum “Battista Grassi”, Sapienza University of Rome. MGP: Museum of Geology and Paleontology of Turin. MPCCSM: Museum of Pre-Classical Civilizations of Southern Murgia. MPUN: Paleontological Museum of the University of Naples Federico II. MUCIV: Archaeozoology laboratory of the Museum of Civilizations (includes the National Prehistoric Ethnographic Museum “Luigi Pigorini”). MUST: University Museum of Earth Sciences, Sapienza University of Rome (includes the Museum of Paleontology, MPUR). PF: PaleoFactory Laboratory, Sapienza University of Rome. SMCC: Speleological Museum of Castellana Caves “Franco Anelli”.

## Results and discussion

## Morphometry

Suidae from the late Middle-Late Pleistocene of Apulia display a huge dental variability (Fig. 2, 3, and 4). Nevertheless, the statistical analysis of M<sub>3</sub>L supports a significant separation between a large-sized interglacial group and a small-sized glacial + Early Holocene one. In fact, the MIS 1 reference sample is classified as distinct from the interglacial samples (MIS 3 and MIS 5), Melpignano, and the extant Italian population, but not from glacial groups (MIS 2 and MIS 4) (Fig. 5).

Lower third molars are larger in Melpignano ( $M_3L$  40.5 mm) and interglacial specimens (Avetrana 2–7:  $M_3L$  mean 41.1 mm, min 40.3 mm, max 41.9 mm; Avetrana 8:  $M_3L$  mean 41.1 mm, min 40.0 mm, max 42.2 mm) than in the glacial ones (the longest specimen is from Grotta Paglicci:  $M_3L$  38.8 mm) and extant wild boar ( $M_3L$  mean 38.2 mm, min 34.0 mm, max 42.5 mm) (Tab. 4, Fig. 4). These values significantly exceed the 2–7% range of sex-related differences observed in previous studies (Made, 1991; Kuşatman, 1991). Indeed, the average interglacial (MIS 3 and MIS 5)  $M_3L$  of 41.1 mm is 20.2% larger than the average glacial length of 32.8 mm, with no overlap in measurement (Tab. 4, Fig. 5). Judging by their size, interglacial suids from Apulia are closer to the Italian Middle Pleistocene remains than to the others Apulian Late Pleistocene ones. Particularly,  $M_3L$  from Avetrana are comparable to Bristie 1 (Trieste) ( $M_3L$  mean 42.3 mm, min 39.6 mm, max 44.1 mm;  $M_3W$  mean 18.9 mm, min 18.0 mm, max 19.7 mm), Cerè ( $M_3L$  41.5.0 mm,  $M_3W$  17.5 mm) and the upper levels of Torre del Pagliaccetto (Rome) (Id. n.399:  $M_3L$  47.0 mm,  $M_3W$  20.0 mm; Id. n.624:  $M_3L$  41.0 mm  $M_3W$  19.0 mm) chronologically referred to the Middle Pleistocene (Ghezzo et al., 2014; Lugli and Sala, 2000; Caloi and Palombo, 1978; Fabiani, 1919). Conversely, small-sized *S. scrofa* is reported from Grotta della Cala (Salerno) levels dated to  $26.380 \pm 260$   $^{14}\text{C}$  yr BP and  $26.680 \pm 320$   $^{14}\text{C}$  yr BP (MIS 2) ( $M^3L$  30.1 mm,  $M^3W$  17.6 mm;  $M_2L$  25.2 mm,  $M_2W$  17.0 mm;  $M_3L$  27.5 mm,  $M_3W$  15.6 mm) (Boscato et al., 1997). Second molar measurements are consistent with these results (interglacial-glacial divergence:  $M^2L$  3.2%,  $M_2L$  9.24%), with late Middle Pleistocene and interglacial specimens of larger size than the rest of the fossil material, whereas first molars do not evidence a clear pattern (Tab. 3 and 4, Fig. 3 and 4).

The postcranial sample is too limited to draw significant considerations, nevertheless interglacial specimens from Aventrana are larger than

**Table 4** – Measurements (mm) of the lower teeth of *Sus scrofa* from the late Middle Pleistocene-Early Holocene sites of Apulia and the extant Italian wild boar.

Site	Reference	MIS	P <sup>2</sup> L	P <sup>2</sup> W	P <sup>3</sup> L	P <sup>3</sup> W	P <sup>4</sup> L	P <sup>4</sup> W	M <sup>1</sup> L	M <sup>1</sup> W	M <sup>2</sup> L	M <sup>2</sup> W	M <sup>3</sup> L	M <sup>3</sup> W
Melpignano	This work	99-8					13.8	8.5	18.3	10.8	21.5	15.1	40.5	16.4
Grotta delle Striare	This work	27-5							(min=16.8- max=19.2)	(min=9.2- max=11.7)				
Avetrana 2-7	This work	5					16.2	11.8	25.9	17.5	41.1	17.7	(min=10.3- max=11.9)	(min=17.5- max=17.9)
Grotta dei Giganti	This work	4												
Grotta Zinzulusa - B6	This work	4	13.2	7.6			15.3	9.0	15.6	10.6	22.8	15.1	37.1	16.7
Grotta Mario Bernardini	This work	4												
Grotta Uluzzo C	This work	4												
Grotta Laceduza	This work	4												
Avetrana 8	This work	3	12.2	4.7	14.6	6.7	15.6	9.4	17.3	12.7	21.2	12.7	30.5	14.6
Fondo Focene	This work	2	11.6	6.8	(min=14.4- max=14.8)	(min=6.4- max=7.1)	(min=15.1- max=16.1)	(min=9.0- max=9.9)	(min=17.1- max=17.5)	(min=11.9- max=13.7)	23.8	16.6	41.1	17.9
Grotta Zinzulusa - B5-3	This work	2												
Grotta Paglicci	Boscato, 1994	2												
SMA-esterno - Epigravettian	This work	2												
Grotta Romanelli	This work	1												
Grotta della Jena	This work	1												
Grotta delle Mura	Bon and Boscato, 1993	1												
Grotta delle Pazzicche	This work	1												
Extant <i>Sus scrofa</i>	Kuşatman, 1991; This work	11.5	5.3	14.2	7.4	14.7	9.3	15.5	11.9	20.7	14.9	38.2	17.0	

**Table 5** – Measurements (mm) of postcranial remains and associated body mass estimates (Kg) of *Sus scrofa* from the late Middle Pleistocene-Early Holocene sites of Apulia.

Site	Reference	MIS	Humerus Bd	Humerus estimate	Radius Bd	Radius estimate	Talus GLI	Talus estimate
Melpignano	This work	?9–8			37.4	111.4		
Grotta Mario Bernardini	This work	4	47.7	159.1				
Avetrana 8	This work	3	51.2 (min=50.4- max=52)	198.2 (min=188.7- max=207.7)			58.9 (min=57.1- max=60.8)	289.6 (min=261.3- max=317.9)
Fondo Focone	This work	2			34.3	89.8		
Grotta Paglicci	Boscato, 1994	2			41.7	145.7	49.2	166.3
Grotta Romanelli	This work	1	45.3	137.1				

the other specimens (Humerus Bd: 50.4 mm, 52 mm; Talus length: 57.1 mm; 60.8 mm) and close to the Middle Pleistocene Bristie 1 (Humerus Bd: 49.5 mm; 52.3 mm) or even larger (Talus length: 51 mm; 54 mm; 57.5 mm) (Tab. 5). Moreover, at least for the talus, glacial-interglacial differences (16.5%) slightly exceed the range of sexual variability of modern wild boar (Kuşatman, 1991).

The large size of Middle Pleistocene European wild boar is widely recognized among researchers (Lister et al., 2010; Guérin and Faure, 1997), but no agreement exists on the time and mode of the size reduction. Our results reveal that in the late Middle-Late Pleistocene there is no progressive trend towards small dimensions, as assumed in previous studies (e.g. Guérin and Faure, 1997; Faure and Guérin, 1983; Hünermann, 1975, 1969). Conversely, we observe several shifts in size, related to interglacial-glacial cycles. The only apparent discrepancy is the interglacial but small-sized MIS 1 sample (Grotta Romanelli, Grotta della Jena, Grotta delle Mura, Grotta delle Prazziche). However, the Early Holocene was characterized by a cold and dry environment (Giustini et al., 2018; Martrat et al., 2014; Sadori and Narcisi, 2001), hence this exception actually proves the general pattern observed in this study (smaller size in colder settings). The sample from MIS 2 displays wide amplitude of  $M_3L$  (min 28.2 mm; max 38.8 mm). The variability of this group, unfortunately represented by only two specimens (Tab. 4), exceeds that of the extant sample, even if it does not overlap with MIS 3 and MIS 5 specimens.

## Body mass

Predicted values vary greatly between different measured elements (teeth:  $M^1L$  min estimate: 49.4 kg;  $M_3L$  max estimate: 127 kg; Humerus Bd min estimate: 137.1 kg; max estimate: 207.7 kg; Tab. 5 and 6). Therefore, it is not possible to use body mass as a common currency to compare different morphometric measurements. Suidae are generalist mammals that retain primitive anatomical features (e.g. bunodont dentition, four-toes foot), and the difficulties in reconstructing the body mass of this group of animals are well known in the literature (Fortelius et al., 1996; Janis, 1990; Scott, 1990). For the same reason, we would have expected more reliable predictions based on the talus length, being the equations provided by Tsubamoto (2014) developed with the aim of reconstruct the body mass of Palaeogene mammals with difficult systematics affinities. However, this is not the case either. In fact, talus-based results are clearly overestimated (Avetrana max estimate: 317.9 kg), predicting body masses comparable only to the largest living wild boars from the Russian Far East (Heptner et al., 1988).

Comparing the outcomes obtained for the extant Italian sample with values directly measured on living animals — between 58.9 Kg and 85.6 Kg (Apollonio, 2003) — it can be supported that second molars provide the most reliable results — being between 56.9 kg and 75.3 kg — in agreement with previous studies (Fortelius et al., 1996; Janis, 1990). For what concerns the other teeth, first and third molars estimates are quite divergent. Precisely, first molars underestimate the individual body mass, whereas third molars overestimate it. Only the latter case has an explanation from the literature, as a limitation of using

ungulate-fitted equations for omnivores, which possess proportionally longer teeth (Janis, 1990).

According to second molar predictions, the heaviest wild boars occurred during interglacial stages, with estimated mean values in each case above the ones predicted for the glacial and extant samples (Tab. 4). In fact,  $M^2L$  predicts an average body mass of 76.3 kg for the interglacial sample (MIS 5 and MIS 3), whereas average values are 73.9 kg for the glacial one (MIS 4 and MIS 2) and 65.7 kg for the extant population.  $M_2L$  provides similar results, 75.5 kg during interglacial periods, whereas average glacial, early Holocene and extant values are respectively 68.7 kg, 72.2 kg, and 64.5 kg.  $M_1L$ ,  $M^3L$ , and  $M_3L$  estimates are consistent with these results, even though they provide less reliable absolute values. The only exception are  $M^1L$  estimates, which are biased, however, by the poor representativeness of the first upper molar in the interglacial sample (only one specimen) and by the high percentage of young individuals — thus with relatively longer teeth, being less affected by wear — in the Early Holocene sample.

## Ecomorphological adaptations

Many generalist, wide-ranging large mammal species, display several ecomorphological adaptations in accordance with the environmental variations encountered throughout their geographic range. The extant wild boar roughly follows Bergmann's ecogeographical rule (Bergmann, 1847), that predicts larger size in colder climates (Albarella et al., 2009; Faure and Guérin, 1983). This pattern has been suggested to be valid even through time for other species, with large-sized *Crocuta crocuta* (Klein and Scott, 1989), *Bos primigenius* (Wright, 2013; Cerilli and Petronio, 1992) and *Canis lupus* (Mecozzi and Bartolini Lucenti, 2018; Sansalone et al., 2015) reported from glacial stages. However, in fossil *S. scrofa* from Apulia large dimensions are, conversely, associated with interglacial stages, and this study focuses on a well-constrained region, hence excluding a geographical-driven explanation for the observed values.

Moreover, several studies have casted doubt on the generality of Bergmann's rule (e.g. Huston and Wolverton, 2011; McNab, 2010; Meiri et al., 2007; Yom-Tov and Geffen, 2006; Ashton et al., 2000; Dayan et al., 1991; Demment and Van Soest, 1985). Indeed, contrary to the rule predictions, several fossil mammals underwent drops in size in colder climatic conditions (Lister et al., 2010; Collinge, 2001; Kurten, 1968) or exhibited a series of size fluctuations for which climate is hardly the solely responsible (Marciszak et al., 2019; Made et al., 2014). Asperen (2010) examined late Middle Pleistocene caballoid horses from Europe and concluded that the glacial specimens were in general smaller than the interglacial ones, interpreting this evidence as the result of an adaptive response to the open-environments degradation and the consequently lower environmental productivity. Similarly, Mazza and Bertini (2013) observed evidence of climate-driven body size changes in Pleistocene hippopotamuses, with larger specimens found in warmer intervals and smaller individuals in colder settings. The authors argued for the combined importance of the short-term Pleistocene climatic cycling (recurrent stresses) and the long-term

**Table 6** – Body mass estimates (Kg) derived from the molars of *Sus scrofa* from the late Middle Pleistocene-Early Holocene sites of Apulia and the extant Italian wild boar.

Site	Reference	MIS	M <sup>1</sup> L	M <sup>2</sup> L	M <sup>3</sup> L	M <sub>1</sub> L	M <sub>2</sub> L	M <sub>3</sub> L
Melpignano	This work	?9-8			109.5	58.1	67.1	121.9
Grotta delle Striare	This work	?7-5				(min=53.5- max=60.9)		
Grotta delle Tre Porte	This work	?7-5		106.4		51.6		
Avetrana 2-7	This work	5			107.5 (min=105.5- max=109.8)		80.6	123.7 (min=121.3- max=126.1)
Grotta dei Giganti	This work	4					71.1	111.7
Grotta Zinzulusa - B6	This work	4	54.0			49.8	68.9	99.3 (min=98.7- max=99.8)
Grotta Mario Bernardini	This work	4				52.9		
Grotta Uluzzo C	This work	4						87.8
Grotta Laceduzza	This work	4					66.1	92.0
Avetrana 8	This work	3	49.4	76.3 (min=73.8- max=78.9)	105.3 (min=98.2- max=111.2)	55.0 (min=54.4- max=55.7)	74.1 (min=68.3- max=79.0)	123.7 (min=120.4- max=127.0)
Fondo Focone	This work	2		69.5				
Grotta Paglicci	Boscato, 1994	2						116.8
Grotta Zinzulusa - B5-3	This work	2				53.2 (min=51.6- max=54.4)		
SMA-esterno - Gravettian	This work	2			78.3			
SMA-esterno - Epigravettian	This work	2						85.1
Grotta Romanelli	This work	1	59.6			53.2 (min=51.6- max=54.7)	71.3	103.0 (min=96.5- max=109.6)
Grotta della Jena	This work	1						87.5
Grotta delle Mura	Bon and Boscato, 1993	1			96.5		73.8	102.5
Grotta delle Prazziche	This work	1	55.9 (min=54.0- max=58.7)	66.2	94.9 (min=92.6- max=97.4)	56.6	71.4	108.1 (min=107.5- max=109.0)
Extant <i>Sus scrofa</i>	Kuşatman, 1991; This work		51.7 (min=47.5- max=58.7)	65.7	101.4 (min=95.1- max=105.3)	49.4 (min=43.2- max=57.2)	64.5 (min=56.9- max=74.1)	114.9 (min=102.5- max=127.9)

Quaternary habitat fragmentation (general deterioration), in promoting less favourable environmental conditions.

During glacial periods, in the Mediterranean area the reduced availability of trophic resources was pronounced, with generally open palaeoenvironments dominant (steppe and/or grasslands) (Renault-Miskovsky et al., 2015, 2011; Allen and Huntley, 2009; among others). Moreover, Apulia is a relatively isolated peninsula, whose geographical setting may have exacerbated the patchiness of the environment in cooler and drier stages. Therefore, a great deterioration and fragmentation of the forest, and consequently the decrease of the available trophic resources and the increase of the competition, might have promoted the observed size reductions in *S. scrofa* during the glacial stages.

Indeed, in the low trophic Mediterranean area, extant wild boars from Andalusia (southern Spain), Corsica (France), and Sardinia (Italy) are usually grouped in the subspecies *S. scrofa meridionalis*, whose small size is the most distinctive trait (Groves, 2007). However, Corsica and Sardinia are islands whereas Andalusia is a low trophic, but continental, region, similarly to Apulia.

These examples testify that “rules” alternative to Bergmann’s thermoregulatory principle better explain the wild boar geographic size pattern, such as the “resource rule” (resource availability; McNab, 2010) or the “eNPP rule” (amount of ecologically and evolutionarily relevant net primary productivity; Huston and Wolverton, 2011). Indeed, several empirical studies that have evaluated the adherence of different species to Bergmann’s principle (Freckleton et al., 2003; Meiri and Dayan, 2003; Ashton et al., 2000), have also denoted a dissimilar at-

titude to follow the rule in large and small mammals, with no or different patterns more common in smaller species (e.g. adherence to the rule: >85% in species larger than 5 kg; <50% in species smaller than 0.5 kg; Meiri and Dayan, 2003). However, on a theoretical ground this differential response is in contrast with Bergmann’s thermoregulatory explanation, since small mammals are supposed to display a stricter adherence to the rule, being affected by a higher rate of heat dispersion (Ashton et al., 2000; Steudel et al., 1994). The paradox is solved if another underlying principle is proposed to describe the observed pattern: the availability of trophic resources (Huston and Wolverton, 2011; McNab, 2010). In fact, in this case the differences between large and small mammals can be interpreted as differences between, respectively, K- and r-selected species (which differ in terms of productivity and efficiency of reproduction), the latter being more prone to evolve faster adaptations to cope with resources oscillations (Tuomi, 1980; Pianka, 1970). Palaeontologists are well aware of the dissimilar rate of change in small and large mammals, which enhance the importance of integrated biochronological scales (Masini and Sala, 2007; Gliozzi et al., 1997).

Of course, the dichotomy between r and K strategy is an empirical generalization and species are distributed along an ecological continuum (Tuomi, 1980; Pianka, 1970). Indeed, despite being large mammals, suids are relatively r-selected species (Martínez-Navarro et al., 2015) and this is reflected in the great biochronological value of this group (e.g. White and Harris, 1977). The wild boar, in particular, can be considered the most r-selected extant large mammal, being the un-

**Table 7** – Regression equations used for estimating the body mass of *Sus scrofa* from the late Middle Pleistocene-Early Holocene sites of Apulia and the extant Italian wild boar.

Equation	Reference	Measure (a)	Slope (b)	Intercept (c)
Body mass=axb+c	Damuth, 1990	M <sup>1</sup> L	3.09	1.21
Body mass=axb+c	Damuth, 1990	M <sup>2</sup> L	3.03	1.06
Body mass=axb+c	Damuth, 1990	M <sup>3</sup> L	2.81	1.29
Body mass=axb+c	Damuth, 1990	M <sub>1</sub> L	3.11	1.24
Body mass=axb+c	Damuth, 1990	M <sub>2</sub> L	3.07	1.07
Body mass=axb+c	Damuth, 1990	M <sub>3</sub> L	2.99	0.80
Log <sub>10</sub> (Body mass)=Log <sub>10</sub> (a)xb+c	Scott, 1990	Humerus Bd	3.09	0.02
Log <sub>10</sub> (Body mass)=Log <sub>10</sub> (a)xb+c	Scott, 1990	Radius Bd	2.48	1.14
(Ln(Body mass)=Ln(a)xb+c)x1.345	Tsubamoto, 2014	Talus GLI	3.12	-0.46

gulate species with the highest reproductive potential in relation to its body mass (Bieber and Ruf, 2005).

This perspective provides further support for the evolutionary scenario reconstructed for the late Middle Pleistocene-Early Holocene *S. scrofa* in Apulia and challenges the supposed conformity of the species to Bergmann's rule. In fact, the ecological strategy adopted by the species is in contrast with an adherence to the rule, but explains how the wild boar managed to cope with the severe resource fluctuations occurred during glacial stages with a size reduction pattern, whereas other large mammals reacted by modifying their range or went extinct (see online resource Supplemental S1).

## Conclusion

The wild boar is one of the most common species in the Middle-Late Pleistocene fossil record of Italy and Europe. Nevertheless, most of the material has never been described or figured, as in the case of several Italian localities assuming a key role for the Italian large mammal biochronological scale, such as Ponte Galeria, Isernia La Pineta, and Fontana Ranuccio.

The fossils from several late Middle Pleistocene to Early Holocene sites from the Apulian Peninsula are here considered for the first time. Our data give no support to the hypothesis of a progressive trend towards small dimension, as sometimes assumed in previous studies. Conversely, morphometric comparisons, body mass estimates and statistical analysis identify two distinct groups of wild boars: large-sized from interglacial stages; small-sized from glacial ones and the cold Early Holocene. The interglacial group is 20.2% larger in M<sub>3</sub>L than both glacial and Early Holocene specimens, with no overlap in measurements. Moreover, this divergence markedly exceeds the range of sexual size dimorphism in extant wild boars (2–7%). M<sup>1</sup>L, M<sub>2</sub>L, and M<sup>3</sup>L are consistent with these results, even though differences are less pronounced for second molars and we lack M<sup>3</sup> from glacial sites. First molars and postcranial remains are too limited to draw significant considerations.

In the first cold Late Pleistocene stage, the straight-tusked elephant and hippo disappeared from Italy. This clearly indicates an early climatic recrudescence, which affected the composition of the large mammal fauna. The results of this work suggest that the wild boar apparently responded to strong climatic fluctuations with a size reduction pattern, whereas other large mammals reacted by modifying their range or went extinct. Moreover, climate could have played an indirect role in leading to the onset of a typical Mediterranean environment, reducing the availability of trophic resources. This pattern is in contrast with the supposed adherence of *S. scrofa* to Bergmann's rule, but in agreement with the r-strategy adopted by the species.

Finally, in order to provide a more articulated framework of the climate-terrestrial ecosystems interactions during the late Quaternary, the comparison to Peninsular Italy and other region of the Euro-Mediterranean area is needed. 

## References

- Abbazzi L., Fanfani F., Ferretti M.P., Rook L., Cattani L., Masini F., Mallegni F., Negrino F., Tozzi C., 2000. New Human Remains of Archaic *Homo sapiens* and Lower Palaeolithic Industries from Visogliano (Duino Aurisina, Trieste, Italy). *J. Archaeol. Sci.* 27: 1173–1186.
- Albarella U., Dobney K., Rowley-Conwy, P. 2009. Size and shape of the Eurasian wild boar (*Sus scrofa*), with a view to the reconstruction of its Holocene history. *Environ. Archaeol.* 14: 103–136.
- Allen J.R., Huntley B., 2009. Last Interglacial palaeovegetation, palaeoenvironments and chronology: a new record from Lago Grande di Monticchio, southern Italy. *Quat. Sci. Rev.* 28: 1521–1538.
- Ambrosetti P., Bartolomei G., De Giuli C., Ficcarelli G., Torre D., 1979. La breccia ossifera di Silvia (Aurisina-Sistiana) nel corso di Trieste. *Boll. Soc. Paleontolog. Ital.* 18: 207–220. [in Italian]
- Anelli F., 1956. Un raro reperto di stambocco nella grotta della Jena presso Castellana (Bari). 7 Congresso Nazionale Di Speleologia. 105–107. [in Italian]
- Anzidei A.P., Bulgarelli G.M., Catalano P., Cerilli E., Gallotti R., Lemorini C., Milli S., Palombo M.R., Pantano W., Santucci E., 2012. Ongoing research at the late Middle Pleistocene site of La Polledrara di Cecanibbio (central Italy), with emphasis on human-elephant relationships. *Quat. Int.* 255: 171–187.
- Apollonio M., 2003. *Sus scrofa* (Linnaeus, 1758), in Boitani L., Lovari S., Vigna Taglianti A. (Eds.) *Fauna d'Italia. Mammalia III. Carnivora-Artiodactyla*. Calderini, Bologna. 261–271. [in Italian]
- Ashton K.G., Tracy M.C., de Queiroz A., 2000. Is Bergmann's Rule Valid for Mammals? *Am. Nat.* 156: 390–415.
- Asperen E.N. van, 2010. Ecomorphological adaptations to climate and substrate in late Middle Pleistocene caballoid horses. *Palaeogeogr., Palaeoclimatol., Palaeoecol.* 297: 584–596.
- Aureli D., Ronchitelli A., 2018. The Lower Tyrrhenian Versant: was it a techno-cultural area during the Middle Palaeolithic? Evolution of the lithic industries of the Riparo del Molare sequence in the frame of Neanderthal peopling dynamics in Italy, in: Borgia V., Cristiani E. (Eds.) *Palaeolithic Italy Advanced Studies on Early Human Adaptations in the Apennine Peninsula*. Sidestone Press Academics, Leiden. 59–94.
- Azzaroli A., 1977. The Villafranchian stage in Italy and the Plio-Pleistocene boundary. *G. Geol.* 41: 61–79.
- Baillié H., 2015. Les industries Lithiques des occupations Épigravettiennes de SMA-Esterno (fouilles 2007–2010) (Santa Maria di Agnano, Ostuni, Italie). *Bulletin du Musée d'Anthropologie préhistorique de Monaco* 55: 105–122. [in French]
- Ballari S.A., Barrios-Garcia M.N., 2014. A review of wild boar *Sus scrofa* diet and factors affecting food selection in native and introduced ranges. *Mamm. Rev.* 44: 124–134.
- Bedetti C., Palombo M.R., Sardella R., 2001. Last occurrences of large mammals and birds in the Late Quaternary of the Italian peninsula. *La Terra Degli Elefanti – The World of Elephants*, Rome. 701–703.
- Bellucci L., Sardella R., Rook L., 2015. Large mammal biochronology framework in Europe at Jaramillo: the Epivilafranchian as a formal biochron. *Quat. Int.* 389: 84–89.
- Bergmann C., 1847. Über die Verhältnisse der Wärmeökonomie der Thiere zu ihrer Grösse. *Göttinger Studien*. [in German]
- Berto C., Boscatto P., Boschin F., Luzi E., Ronchitelli A., 2017. Paleoenvironmental and paleoclimatic context during the Upper Palaeolithic (late Upper Pleistocene) in the Italian Peninsula. The small mammal record from Grotta Paglicci (Rignano Garganico, Foggia, Southern Italy). *Quat. Sci. Rev.* 168: 30–41.
- Bieber C., Ruf T., 2005. Population dynamics in wild boar *Sus scrofa*: ecology, elasticity of growth rate and implications for the management of pulsed resource consumers. *J. Appl. Ecol.* 42: 1203–1213.
- Blanc A.C., 1958. Industria musteriana su calcare e su valve di Meretrix chione associata con fossili di Elefante e Rinoceronte, in nuovi giacimenti costieri del Capo di Leuca. *Quaternaria* 5: 308–313. [in Italian]
- Blanc A.C., 1962. Relazione sulle osservazioni e ricerche stratigrafiche eseguite nella Grotta Zinzulusa (Castro Marina, Lecce) e sull'esito di alcuni sopralluoghi nel Salento. *Quaternaria* 5: 330–334. [in Italian]
- Blanc G.A., 1920. Grotta Romanelli I. Stratigrafia dei depositi e natura e origine di essi. *Archivio per l'Antropologia e la Etnologia* 50: 1–39. [in Italian]
- Bologna P., 1992. La fauna Pleistocenica di Cursi (Lecce). M.Sc. thesis, Sapienza Università di Roma, Roma. [in Italian]
- Bologna P., Di Stefano G., Manzi G., Petronio C., Sardella R., Squazzini E., 1994. Late Pleistocene mammals from the Melpignano (LE) “Ventarole”: preliminary analysis and correlations. *Boll. Soc. Paleontolog. Ital.* 33: 265–274.
- Bon M., Boscatto P., 1993. Analisi paleontologica e paleoecologica di macro e micromamiferi dei livelli romanelliani e mesolitici della Grotta delle Mura (Monopoli, Bari). *Quaternaria Nova* 3: 53–104. [in Italian]
- Borzatti von Löwenstern E., 1966. Il Neolitico di grotta delle Prazziche. Atti della X Riunione Scientifica dell'Istituto Italiano di Preistoria e Protostoria. 7–12. [in Italian]
- Borzatti von Löwenstern E., 1969. Industrie romanelliane e neolitiche nella grotta delle Prazziche (Novaglie-Lecce). *Riv. Sci. Preist.* 24: 91–143. [in Italian]
- Borzatti von Löwenstern E., Magaldi D., 1969. Risultati conclusivi dello studio paleontologico e sedimentologico della grotta di Uluzzo C (Nardò, Lecce). *Riv. Sci. Preist.* 24: 15–64. [in Italian]

- Boscato P., 1994. Grotta Paglicci: la fauna a grandi mammiferi degli strati 22–24 (Gravettiano antico-Aurignaziano). Riv. Sci. Preist. 46: 145–176. [in Italian]
- Boscato P., Ronchitelli A., Wierer U., 1997. Il Gravettiano antico della Grotta della Cala a Marina di Camerota: paletnologia e ambiente. Riv. Sci. Preist. 48: 97–186. [in Italian]
- Botti U., 1874a. La Zinzolosa. Tipografia di G. Barbera. [in Italian]
- Botti U., 1874b. Sulla scoperta di ossa fossili in Terra d'Otranto. Boll. Regio Comitato Geol. 1: 7–8. [in Italian]
- Botti U., 1890. La grotta ossifera di Cardamone in Terra d'Otranto. Boll. Soc. Geol. Ital. 9: 659–716. [in Italian]
- Braun I.M., Palombo M.R., 2012. *Mammuthus primigenius* in the cave and portable art: An overview with a short account on the elephant fossil record in Southern Europe during the last glacial. Quat. Int. 276–277: 61–76.
- Calcagnile L., Sardella R., Mazzini I., Giustini F., Brilli M., D'Elia M., Braione E., Conti J., Mecozzi B., Bona F., Iurino D.A., Lembo G., Muttilo B., Quarta G., 2019. New radiocarbon dating results from the upper Paleolithic–Mesolithic levels in Grotta Romanelli (Apulia, Southern Italy). Radiocarbon 1–10.
- Caloi L., Palombo M.R., 1978. Anfibi, rettili e mammiferi di Torre del Pagliacetto (Torre in Pietra, Roma). Quaternaria 20: 315–428. [in Italian]
- Caloi L., Palombo M.R., 1979. Resti di mammiferi del Pleistocene Medio di Malagrotta (Roma). Boll. Serv. Geol. Ital. 100: 141–188. [in Italian]
- Cancellieri L., 2017. Il sito di Fondo Focene-Trincea B (Ugento, Lecce) nel contesto dell'Epigravettiano antico dell'Italia peninsulare orientale. Studi di Preistoria e Protostoria 4: 79–86. [in Italian]
- Cardini L., 1962. Le faune dei nuovi orizzonti della Grotta Zinzulosa. Quaternaria 5: 334. [in Italian]
- Cardini L., 1965. Sui giacimenti paleolitici di recente scoperti nella penisola Salentina. VI Congresso Internazionale Di Scienze Preistoriche e Protostoriche, Florence. 144–148. [in Italian]
- Cerilli E., Petronio C., 1992. Biometrical variations of *Bos primigenius* Bojanus. Proceedings of the International Symposium "Ongules/Ungulates 91", Toulouse. 37–42.
- Cherin M., Sorbelli L., Crotti M., Iurino D.A., Sardella R., Souron A., 2018. New material of *Sus strozzii* (Suidae, Mammalia) from the Early Pleistocene of Italy and a phylogenetic analysis of suines. Quat. Sci. Rev. 194: 94–115.
- Cigna A., Orlandi G., 1954. La grotta delle Striare, caverna ossifera della costa jonica (Lecce). VI Congresso Nazionale Di Speleologia, Trieste. 1–5. [in Italian]
- Collinge S.E., 2001. Body Size and Community Structure in British Pleistocene Mammals. PhD thesis, University College London, London.
- Coppola D., Pinto N. De, Pellegrino M., Baills H., Stasolla V., Dewailly M., Féret S., Breichner H., Russo J., Bartolomeo J., 2017. Santa Maria di Agnano (Ostuni, Puglia). Chronique des activités archéologiques de l'École française de Rome. [in French]
- Corridi C., 1989. La fauna di Fondo Cattie (Maglie) nel quadro biocronologico del Pleistocene superiore del Salento. Riv. Sci. Preist. 42: 331–337. [in Italian]
- Damuth J., 1990. Problems in estimating body masses of archaic ungulates using dental measurements. In: Damuth J., MacFadden, B.J. (Eds.) Body Size in Mammalian Paleobiology: Estimation and Biological Implications. Cambridge University Press. 229–254.
- Dayan T., Simberloff D., Tchernov E., Yom-Tov Y., 1991. Calibrating the paleothermometer: climate, communities, and the evolution of size. Paleobiology 17: 189–199.
- Del Campana D., 1909. Vertebrati fossili di Monte Tignoso (Livorno). Boll. Soc. Geol. Ital. 28: 349–388. [in Italian]
- Demment M.W., Van Soest P.J., 1985. A Nutritional Explanation for Body-Size Patterns of Ruminant and Nonruminant Herbivores. Am. Nat. 125: 641–672.
- Driesch A. von den, 1976. A guide to the measurement of animal bones from archaeological sites. Peabody Museum Bulletin 1: 1–137.
- Fabiani R., 1919. I mammiferi quaternari della regione Veneta. Mem. Ist. Geol. Univ. 5: 1–174. [in Italian]
- Faure M., Guérin C., 1983. Le *Sus scrofa* (Mammalia, Artiodactyla, Suidae) du gisement pléistocène supérieur de Jaurens à Nespolu, Corrèze, France. Nouvelles Archives du Muséum d'Histoire Naturelle de Lyon 21: 45–63. [in French]
- Fortelius M., van der Made J., Bernor R.L., 1996. Middle and Late Miocene Suoidea of Central Europe and the Eastern Mediterranean: evolution, biogeography, and paleoecology. The evolution of Western Eurasian Neogene mammal faunas. 348–377.
- Frantz L., Meijaard E., Gongora J., Haile J., Groenend M.A.M., Larson G., 2016. The Evolution of Suidae. Annu. Rev. Anim. Biosci. 4: 61–85.
- Fraendorf M., Gethöffer F., Siebert U., Keulung O., 2016. The influence of environmental and physiological factors on the litter size of wild boar (*Sus scrofa*) in an agriculture dominated area in Germany. Sci. Total Environ. 541: 877–882.
- Freckleton R.P., Harvey P.H., Pagel M., 2003. Bergmann's Rule and Body Size in Mammals. Am. Nat. 161: 821–825.
- Gallo-Orsi U., Macchi E., Perrone A., Durio P., 1995. Biometric data and growth rates of a wild boar population living in the Italian Alps. J. Mount. Ecol. 3: 60–63.
- Gaillard J.-M., Pontier D., Allainé D., Lebreton J.D., Trouvilliez J., Clobert J., Allaine D., 1989. An Analysis of Demographic Tactics in Birds and Mammals. Oikos 56: 59–76.
- Genov P.V., 1999. A review of the cranial characteristics of the Wild Boar (*Sus scrofa* Linnaeus 1758), with systematic conclusions. Mamm. Rev. 29: 205–234.
- Ghezzo E., Berté D.F., Sala B., 2014. The revaluation of Galerian Canidae, Felidae and Mustelidae of the Cerè Cave (Verona, Northeastern Italy). Quat. Int. 339: 76–89.
- Giustini F., Bona F., Brilli M., Conti J., D'Agostino A., Lembo G., Mazzini I., Mecozzi B., Muttilo B., Sardella R., 2018. An introduction to the early Holocene elonian deposits of Grotta Romanello, Apulia, Southern Italy. Alp. Mediterr. Quat. 31: 135–139.
- Gliozzi E., Abbazzi L., Argenti P., Azzaroli A., Caloi L., Capasso Barbato L., Di Stefano G., Esu D., Ficcarelli G., Girotti O., Kotsakis T., Masini F., Mazza P., Mezzabotta C., Palombo M.R., Petronio C., Rook L., Sala B., Sardella R., Zanalda E., Torre D., 1997. Biochronology of selected mammals, molluscs and ostracodes from the Middle Pliocene to the Late Pleistocene in Italy. The state of the art. Riv. Ital. Paleontolog. Stratig. 103: 369–388.
- Goldfuss K.A., 1823. Osteologische Beiträge zur Kenntniß verschiedener Säugetiere der Vorwelt. Nova Acta Physico-Medica Academiae Caesareae Leopoldino-Caroliniae Naturae Curiosorum 11: 482–485. [in German]
- Groves C.P., 1981. Ancestors for the pigs: taxonomy and phylogeny of the genus *Sus*. Dept. of Prehistory, Research School of Pacific Studies, Australian National University, Canberra.
- Groves C.P., 2007. Current views on taxonomy and zoogeography of the genus *Sus*. In: Albarella U., Dobney K., Ervynck A., Rowley-Conwy P. (Eds.) Pigs and Humans: 10,000 Years of Interaction. Oxford University Press, New York. 15–29.
- Guérin C., Faure M., 1997. The wild boar (*Sus scrofa priscus*) from the post-Villafranchian Lower Pleistocene of Untermaßfeld. In: Kahlke, R.-D. (Ed.) Das Pleistozän von Untermaßfeld Bei Meiningen (Thüringen), Bonn. 375–384.
- Heptner V.G., Nasimovich A.A., Bannikov A.G., 1988. Mammals of the Soviet Union. Volume I. Amerind Publishing Co., New Delhi.
- Hünermann K.A., 1969. *Sus scrofa priscus* Goldfuss im Pleistozän von Süßenborn bei Weimar. Palaontologische Abhandlungen. Abteilung A, Palaözoologie 3: 611–616. [in German]
- Hünermann K.A., 1975. *Sus scrofa* Linne aus dem Pleistozän von Weimar-Ehringsdorf. Abhandlungen des Zentral Geologischen Instituts 23: 251–263. [in German]
- Huston M.A., Wolverton S., 2011. Regulation of animal size by eNPP, Bergmann's rule, and related phenomena. Ecol. Monogr. 81: 349–405.
- Iannucci A., Gasparik M., Sardella R., 2020. First report of *Sus strozzii* (Suidae, Mammalia) from the Early Pleistocene of Hungary (Dunaalmás) and species distinction based on deciduous teeth. Sci. Nat. 107: 5.
- Janis C.M., 1990. Correlation of cranial and dental variables with body size in ungulates and macropodoids. In: Damuth J., MacFadden, B.J. (Eds.) Body Size in Mammalian Paleobiology: Estimation and Biological Implications. Cambridge University Press. 255–291.
- Klein R.G., Scott K., 1989. Glacial/Interglacial Size Variation in Fossil Spotted Hyenas (*Crocuta crocuta*) from Britain. Quat. Res. 32: 88–95.
- Kurtén B., 1968. Pleistocene Mammals of Europe. Weidenfeld and Nicolson, London.
- Kuşatman B., 1991. The origins of pig domestication with particular reference to the Near East. PhD thesis, University College London, London.
- Lister A.M., Parfitt S.A., Owen F.J., Collinge S.E., Breda M., 2010. Metric analysis of ungulate mammals in the early Middle Pleistocene of Britain, in relation to taxonomy and biostratigraphy: II: Cervidae, Equidae and Suidae. Quat. Int. 228: 157–179.
- Lugli C., Sala B., 2000. La Teriofauna del Pleistocene Medio di Bristie I° (Carso Triestino). Atti del Museo civico di Storia naturale di Trieste 48: 35–58. [in Italian]
- Made J. van der, 1991. Sexual bimodality in some recent pig populations and application of the findings to the study of fossils. Zeitschrift für Säugetierkunde 56: 81–87.
- Made J. van der, Stefanik K., Marciszak A., 2014. The Polish fossil record of the wolf *Canis* and the deer *Alces*, *Capreolus*, *Megaloceros*, *Dama* and *Cervus* in an evolutionary perspective. Quat. Int. 326: 1–25.
- Made J. van der, Rosell J., Blasco R., 2017. Faunas from Atapuerca at the Early–Middle Pleistocene limit: The ungulates from level TD8 in the context of climatic change. Quat. Int. 433: 296–346.
- Marciszak A., Schouwenburg C., Gornig W., Lipecki G., Mackiewicz P., 2019. Morphometric comparison of *Panthera spelaea* (Goldfuss, 1810) from Poland with the lion remains from Eurasia over the last 700 ka. Quat. Sci. Rev. 223: 10590.
- Martínez-Navarro B., Madurell-Malpica J., Ros-Montoya S., Espigares M.-P., Medin T., Hortola P., Palmqvist P., 2015. The Epivilafranchian and the arrival of pigs into Europe. Quat. Int. 389: 131–138.
- Martrat B., Jimenez-Amat P., Zahn R., Grimalt J.O., 2014. Similarities and dissimilarities between the last two deglaciations and interglaciations in the North Atlantic region. Quat. Sci. Rev. 99: 122–134.
- Masini F., Sala B., 2007. Large- and small-mammal distribution patterns and chronostratigraphic boundaries from the Late Pliocene to the Middle Pleistocene of the Italian peninsula. Quat. Int. 160: 43–56.
- Masini F., Sala B., Ambrosetti P., Azzaroli A., Ficcarelli G., Kotsakis T., Rook L., Torre D., 1990. Mammalian faunas of selected Villafranchian and Galerian localities. INQUA SEQS, Subcommission for European Quaternary Stratigraphy, Cromer Symposium, Norwich.
- Maviglia C., 1955. Ossami fossili trovati nella Grotta delle Striare (Otranto). Natura 46: 92–95. [in Italian]
- Mazza P., Bertini A., 2013. Were Pleistocene hippopotamuses exposed to climate driven body size changes? Boreas 42: 194–209.
- McNab B.K., 2010. Geographic and temporal correlations of mammalian size reconsidered: a resource rule. Oecologia 164: 13–23.
- Mecozzi B., Bartolini Lucenti S., 2018. The Late Pleistocene *Canis lupus* (Canidae, Mammalia) from Avetrana (Apulia, Italy): reappraisal and new insights on the European glacial wolves. Ital. J. Geosci. 137: 138–150.
- Mecozzi B., Chakroun A., Baills H., Iurino D.A., Sardella R., Coppola D., 2018. Late Pleistocene mammal faunal assemblage from Cava Spagnulo (Grottaglie, Apulia, SE Italy). Alp. Mediterr. Quat. 31 (1): 5–19.
- Mecozzi B., Bellucci L., Giustini F., Iurino D.A., Mazzini I., Sardella R., 2019a. Large mammal fauna from the late Middle Pleistocene site of Melpignano and San Sidero (Lecce, southern Italy). Paleodays 2019, XIX Riunione annuale SPI, Benevento-Pietrarossa. 24.
- Mecozzi B., Coppola D., Iurino D.A., Sardella R., De Marinis A.M., 2019b. The Late Pleistocene European badger *Meles meles* from Grotta Laceduzza (Brindisi, Apulia, Southern Italy): the analysis of the morphological and biometric variability. Sci. Nat. 106: 13.
- Meiri S., Dayan T., 2003. On the validity of Bergmann's rule. J. Biogeogr. 30: 331–351.
- Meiri S., Yom-Tov Y., Geffen E., 2007. What determines conformity to Bergmann's rule? Global Ecol. Biogeogr. 16: 788–794.
- Minieri M.R., Petronio C., Sardella R., Scarano M., 1995. Le faune a mammiferi del Pleistocene Superiore dell'Italia peninsulare. Atti 1° Convegno Nazionale Di Archeozoologia, Padusa Quadreria 1: 75–87. [in Italian]
- Mirigliano G., 1942. La Cala delle Ossa, presso Palinuro (Salerno). Boll. Soc. Naturalisti in Napoli 54: 1–36. [in Italian]
- Moigne A.-M., Palombo M.R., Belda V., Heriech-Briki D., Kacimi S., Lacombat F., de Lumley M.-A., Moutoussamy J., Rivals F., Quiles J., Testu A., 2006. Les faunes de grands mammifères de la Caune de l'Arago (Tautavel) dans le cadre biochronologique des faunes du Pléistocene moyen italien. L'Anthropologie 110: 788–831. [in French]
- Nores C., Llaneza L., Álvarez Á., 2008. Wild boar *Sus scrofa* mortality by hunting and wolf *Canis lupus* predation: an example in northern Spain. Wildl. Biol. 14: 44–51.
- Okarma H., Jędrzejewska B., Jędrzejewski W., Krasiński Z.A., Milkowski L., 1995. The roles of predation, snow cover, acorn crop, and man-related factors on ungulate mortality in Białowieża Primeval Forest, Poland. Acta Theriol. 40: 197–217.
- Pandolfi L., Petronio C., 2015. A brief review of the occurrences of Pleistocene *Hippopotamus* (Mammalia, Hippopotamidae) in Italy. Geol. Croat. 68: 313–319.

- Pandolfi L., Petronio C., Salari L., 2011. *Bos primigenius* Bojanus, 1827 from the early Late Pleistocene deposit of Avetrana (Southern Italy) and the variation in size of the species in Southern Europe: preliminary report. *J. Geol. Res.* 2011: 1–11.
- Pandolfi L., Petronio C., Salari L., 2013. Catastrophic death assemblages from the Late Pleistocene of Italy: the case of Avetrana karst filling (Taranto, Southern Italy). *Riv. Ital. Paleontol. Stratigr.* 119 (1): 109–124.
- Petronio C., Bellardini F., Arzarello M., Bedetti C., Bellucci L., Cipullo A., Di Stefano G., Pandolfi L., Pavia M., Petrucci M., Sardella R., Salari L., 2008. The deposit of the Late Pleistocene from Avetrana (Taranto, southern Italy): biochronology and palaeoecology. *Il Quaternario* 21: 409–422.
- Petronio C., Bellucci L., Martinetto E., Pandolfi L., Salari L., 2011. Biochronology and paleoenvironmental changes from the Middle Pliocene to the Late Pleistocene in Central Italy. Biochronology and paleoenvironmental changes from the Middle Pliocene to the Late Pleistocene in Central Italy. *Geodiversitas* 33: 485–517.
- Petronio C., Di Canzio E., Salari L., 2007. The Late Pleistocene and Holocene Mammals in Italy: new biochronological and paleoenvironmental data. *Palaeontographica, Abt. A* 279: 147–157.
- Petronio C., Sardella R., 1999. Biochronology of the Pleistocene mammal fauna from Ponte Galeria (Rome) and remarks on the Middle Galerian faunas. *Riv. Ital. Paleontol. Stratigr.* 105: 155–164.
- Pianka E.R., 1970. On r- and K-Selection. *Am. Nat.* 104: 592–597.
- R Core Team, 2019. R language definition. Vienna, Austria: R foundation for statistical computing.
- Renault-Miskovsky J., Baills H., Marquer L., Coppola D., 2015. Santa Maria di Agnano (Ostuni, Brindisi, Italia). Préhistoire et Palynologie, Paléoclimatologie et Paléoenvironnement. *Riv. Sci. Preist.* 65: 5–27. [in French]
- Renault-Miskovsky J., Marquer L., Baills H., Coppola D., 2011. Environnement végétal et paléopédologie du Gravettien et de l'Épigravettien de la Grotte de Santa Maria di Agnano (Ostuni, Brindisi, Italia). Bulletin du Musée d'Anthropologie préhistorique de Monaco 51: 75–86. [in French]
- Romandini M., Cazzanini J., Bortolini E., Boscato P., Boschin F., Carrera L., Nannini N., Tagliacozzo A., Terlato G., Arrighi S., Badino F., Figus C., Lugli F., Mariani G., Oxilia G., Moroni A., Negrino F., Marco P., Riel-Salvatore J., Ronchitelli A., Spinapoli E.E., Benazzi S., 2019. Macromammal and bird assemblages across the Late Middle to Upper Palaeolithic transition in Italy: an extended zooarchaeological review. *Quat. Int.* doi: 10.1016/j.quaint.2019.11.008
- Sadoni L., Narcisi B., 2001. The Postglacial record of environmental history from Lago di Pergusa, Sicily. *The Holocene* 11(6): 655–670.
- Sala B., 1996. Gli animali del giacimento di Isernia La Pineta. In: Peretto C. (Ed.) I reperti paleontologici del giacimento paleolitico di Isernia La Pineta. Cosmo Iannone, Isernia. 41–65. [in Italian]
- Sala B., Masini F., 2007. Late Pliocene and Pleistocene small mammal chronology in the Italian peninsula. *Quat. Int.* 160: 4–16.
- Sala B., Masini F., Ficcarelli G., Rook L., Torre D., 1992. Mammal dispersal events in the Middle and Late Pleistocene of Italy and Western Europe. *Cour. Forschungsinstitut Senckenberg* 153: 59–68.
- Salari L., Petronio C., Kotsakis T., Di Stefano G., Grossi F., Maiorino L., Pandolfi L., Rolfo M.F., Ruiu F.D., Sansalone G., Tagliacozzo A., 2019. Reassessing the faunal assemblages of the Late Pleistocene stratified karst filling from Avetrana (Apulia, southern Italy): the bed 8, palaeoenvironment and biochronology. *Alp. Mediterr. Quat.* 32(2): 1–55.
- Sansalone G., Bertè D.F., Maiorino L., Pandolfi L., 2015. Evolutionary trends and stasis in carnassial teeth of European Pleistocene wolf *Canis lupus* (Mammalia, Canidae). *Quat. Sci. Rev.* 110: 36–48.
- Sardella R., Bedetti C., Bellucci L., Conti N., Coppola D., Di Canzio E., Pavia M., Petronio C., Petrucci M., Salari L., 2005. The Late Pleistocene vertebrate fauna from Avetrana (Taranto, Apulia, Southern Italy): Preliminary report. *Geo.Alp* 2: 25–29.
- Sardella R., Bellucci L., Bona F., Cherin M., Iurino D.A., Rook L., 2018a. Before and after the earliest *Homo* dispersal in Europe: Evidence from the early Pleistocene sites of the Italian Peninsula. *C. R. Palevol* 17: 287–295.
- Sardella R., Mazzini I., Giustini F., Mecozi B., Brilli M., Iurino D.A., Lembo G., Muttillo B., Massussi M., Sigari D., Tucci S., Voltaggio M., 2018b. Grotta Romanelli (Southern Italy, Apulia): legacies and issues in excavating a key site for the Pleistocene of the Mediterranean. *Riv. Ital. Paleontol. Stratigr.* 124: 247–264.
- Sardella R., Iurino D.A., Mecozi B., Sigari D., Bona F., Bellucci L., Coltorti M., Conti J., Lembo G., Muttillo B., Mazzini I., 2019. Grotta Romanelli (Lecce, Southern Italy) Between Past and Future: New Studies and Perspectives for an Archaeo-geosite Symbol of the Palaeolithic in Europe. *Geoheritage*, 11(4): 1413–1432.
- Sarti L., Boscato P., Marto L.M., 1998. Il Musteriano finale di Grotta del Cavallo nel Salento: studio preliminare. Origini: Preistoria e protostoria delle civiltà antiche 22: 45–110. [in Italian]
- Scott K.M., 1990. Postcranial dimensions of ungulates as predictors of body mass. In: Damuth J., MacFadden, B.J. (Eds.) *Body Size in Mammalian Paleobiology: Estimation and Biological Implications*. Cambridge University Press. 301–336.
- Shackleton N.J., 1995. New data on the evolution of pliocene climatic variability. In: Vrba E.S., Denton G.H., Partidge T.C., Burckle L.H. (Eds.) *Paleoclimate and Evolution, with Emphasis on Human Origins*. Yale University Press, New Haven. 242–248.
- Steudel K., Porter W.P., Sher D., 1994. The biophysics of Bergmann's rule: a comparison of the effects of pelage and body size variation on metabolic rate. *Can. J. Zool.* 72: 70–77.
- Strani F., DeMiguel D., Bona F., Sardella R., Bidditti I., Bruni L., De Castro A., Guadagnoli F., Bellucci L., 2018. Ungulate dietary adaptations and palaeoecology of the Middle Pleistocene site of Fontana Ranuccio (Anagni, Central Italy). *Palaeogeogr., Palaeoclimatol., Palaeoecol.* 496: 238–247.
- Tagliacozzo A., 2003. Archeozoologia dei livelli dell'Epigravettiano finale di Grotta Romanelli (Castro, Lecce) strategie di caccia ed economia di sussistenza. In: Fabri P., Ingravallo E., Mangia A. (Eds.) *Grotta Romanelli Nel Centenario Della Sua Scoperta (1900–2000)*. Congedo Editore. 169–216.
- Tsoukala E., Guérin C., 2016. The Rhinocerotidae and Suidae of the Middle Pleistocene from Petralona Cave (Macedonia, Greece). *Acta Zool. Bulg.* 68: 243–264.
- Tsubamoto T., 2014. Estimating body mass from the astragalus in mammals. *Acta Palaeontol. Pol.* 59(2): 259–265.
- Tuomi J., 1980. Mammalian Reproductive Strategies: A Generalized Relation of Litter Size to Body Size. *Oecologia* 45: 39–44.
- White T.D., Harris J.M., 1977. Suid evolution and correlation of African hominid localities. *Science* 198: 13–21.
- Wright E., 2013. The history of the European aurochs (*Bos primigenius*) from the Middle Pleistocene to its extinction: an archaeological investigation of its evolution, morphological variability and response to human exploitation. PhD thesis, University of Sheffield.
- Yom-Tov Y., Geffen E., 2006. Geographic variation in body size: the effects of ambient temperature and precipitation. *Oecologia* 148: 213–218.
- Zanchetta G., Giacco B., Bini M., Sarti L., 2018. Tephrostratigraphy of Grotta del Cavallo, Southern Italy: Insights on the chronology of Middle to Upper Palaeolithic transition in the Mediterranean. *Quat. Sci. Rev.* 182: 65–77.

Associate Editor: P. Raia

## Supplemental information

Additional Supplemental Information may be found in the online version of this article:

**Supplement SI** Late Middle Pleistocene-Early Holocene Apulian fossil Suidae.