



## Short Note

## Distance sampling as a tool for estimating active burrow density in the European ground squirrel (*Spermophilus citellus*): a case study from western Romania

Eliana SEVIANU<sup>1</sup>, Gabriel Bogdan CHISAMERA<sup>2</sup>, Andrada Iulia MARTIN<sup>3</sup>, Ioana COBZARU<sup>2,\*</sup>, Milca PETROVICI<sup>3</sup>

<sup>1</sup>Babeş Bolyai University, Faculty of Environmental Science and Engineering, str. Fântânele 30, 400294, Cluj-Napoca, Romania

<sup>2</sup>Institute of Biology Bucharest, Romanian Academy, 296 Splaiul Independentei, 060031 Bucharest, Romania

<sup>3</sup>West University of Timisoara, Faculty of Chemistry, Biology, Geography, Department of Biology, Pestalozzi 16 street, 300115, Timisoara, Romania

## Keywords:

distance sampling  
European ground squirrel  
burrow openings

## Article history:

Received: 4 March 2025

Accepted: 20 January 2026

## Acknowledgements

I.C. and G.B.C. were funded by project no. RO1567-IBB09/2025 from the Institute of Biology Bucharest of Romanian Academy.

## Abstract

The European ground squirrel (*Spermophilus citellus*), an endangered small mammal endemic to Central and Southeastern Europe, has an unfavourable-inadequate conservation status in Romania. This study assessed the species density of active burrow openings within six areas of the Natura 2000 site ROSAC0108 Lunca Mureşului Inferior in Western Romania. Distance sampling line transect counts of active burrow openings were conducted for two consecutive seasons. Survey design resulted in interannual difference in effective strip width, encounter rate and detection probability. Higher densities were obtained using systematic parallel transects with complete spatial coverage of known colonies than using random line transects in potentially suitable habitat. This study provides insights into methods for estimating population density by using distance sampling approach to counting active burrow openings.

The European ground squirrel (hereafter as EGS; *Spermophilus citellus*) is listed as endangered on the IUCN Red List (Ćosić et al., 2024) and the Red Book of Vertebrates of Romania (Murariu, 2005). It inhabits open short grass steppe, degraded fields with low herbaceous vegetation, earth dikes or dams, and alfalfa crops among cultivated landscapes (Kryštufek, 1999; Kryštufek and Vohralik, 2012; Ramos-Lara et al., 2014; Ružić, 1978). It continues to face population declines despite legal protection (the Bern Convention and the Habitat Directive), and the conservation status remains unfavourable-inadequate, as indicated by two consecutive assessments under Article 17 of the Habitat Directive/Romania (E.E.A., 2025). Assessing the population size is essential for prioritizing appropriate conservation actions (Rammou et al., 2021), although acquiring reliable estimates of wildlife population density through statistically robust sampling techniques remains one of the most significant challenges for field ecologists (Jacquier et al., 2021; Luikart et al., 2010; Thompson, 2004). The species is suitable for population studies using indirect methods like counting active burrow openings (Katona et al., 2002), however, comprehensive and standardized studies in Romania remain limited (Hegyesi et al., 2012), in the absence of a standardised method for estimating EGS abundance across its range (Gedeon et al., 2022; Koshev, 2008).

The main objective of this study was to implement a robust methodology for estimating the density of EGS active burrow openings by employing distance sampling techniques and using two different field sampling designs.

The study was conducted within the Natura 2000 site ROSAC0108 Lunca Mureşului Inferior (western Romania), along the left bank of the Mureş River (90 and 200 m a.s.l.). One of the protected species for which the site was designated is EGS. The site comprises a mosaic of grazed and ungrazed grasslands embedded within an agricultural-forest matrix. Grasslands, representing suitable EGS habitat, cover a total of 2,177 ha, most of which are currently used as pastures, pre-

dominantly grazed by sheep, with local signs of overgrazing and abandonment (MMAP, 2016).


Line transect surveys were conducted in August–September 2022 and June 2023. Each transect was surveyed once by a single observer, and all visible active burrow openings were recorded, measuring the perpendicular distance from the transect line to each detected opening, following standard distance sampling protocols (Buckland et al., 2001). An active burrow opening was identified based on fresh signs of activity, including freshly excavated soil, clear entrances, scrapes at the burrow entrance, fresh droppings and feeding signs (Rammou et al., 2021).

In 2022, surveys targeted known EGS colonies based on past surveys: Sânpetru German (SG), Felnac (FC), and Zădăreni (ZD) (Fig. 1). Within each location, systematic parallel line transects were established, spaced 20 metres apart, corresponding to twice the nominal survey width of the national methodology (Ionescu et al., 2013). The whole areas of the colonies were surveyed, without overlap between adjacent transect detection areas. Individual transect lengths varied depending on local topography, totalling 11.5 km.

In 2023, surveys aimed to estimate the density of active burrow openings across potentially suitable EGS habitat beyond known colonies. Potentially suitable habitat was defined as grazed and ungrazed grasslands characterized by low- to medium-height herbaceous vegetation, excluding crops and forested areas, wetlands, and built-up surfaces. Habitat for each location was delineated based on GPS field data and digitised in ArcGIS to map potentially suitable habitat areas, which were used to guide the placement and spatial extent of random transects. A stratified random sampling design was applied, resulting in 62 line transects with randomly determined starting points and orientations. Habitat surface areas were not used as input parameters in the distance sampling analyses, which were based solely on transect length and perpendicular distance data. Transect lengths varied between 120 and 500 m, covering a total surveyed length of 18.7 km. Fieldwork included broader areas of two of the previously surveyed locations (FC and ZD) and three new: Igriş (IG), Secusigiu (SC), and Munar (MN), where no

\*Corresponding author

Email address: [ioana.cobzaru@ibiol.ro](mailto:ioana.cobzaru@ibiol.ro) (Ioana COBZARU)



**Figure 1** – Location of the six areas where density data of EGS were collected within the Natura 2000 site ROSAC0108 Lunca Mureșului Inferior: IG - Igrîș, SC - Secusigiu, MN - Munar, SG - Sânpetru German, FC - Felnac, ZD - Zădăreni, shown in relation to its geographic distribution according to the IUCN Red List.

prior information on EGS colonies was available. The SG location was excluded as all potential suitable habitat had been surveyed in 2022.

Perpendicular distances to active burrow openings were analysed by means of conventional distance sampling, using Distance 8.0 Release 1 (Thomas et al., 2010). Data were analysed separately for each location within each year, and subsequently pooled at the year level to estimate a single detection function and an annual density estimate. The maximum observed perpendicular distance to an active burrow opening was used as the integration limit for detection function fitting. Detection functions were constrained to be monotone non-increasing with distance, and density estimation was performed without truncation. Several candidate models were fitted, including half-normal and hazard-rate key functions with various adjustment terms. Model selection was based on Akaike’s information criterion (AIC) and the corrected Akaike Information Criterion (AICc), with the model obtaining the lowest score selected as optimal (Burnham and Anderson, 2004). The effective strip width (ESW) for each year was derived from the fitted detection function. Density was calculated by dividing the number of detected active burrow openings by the total effective surveyed area (twice the ESW multiplied by the total transect length), thereby accounting for imperfect detection with distance. The total area covered by transects for each location and year was calculated as twice the maximum observed perpendicular distance from transect to active burrow entrances.

In both years, a half-normal key function with cosine adjustment was selected.

In 2022 a total of 437 active burrow openings were recorded, corresponding to an encounter rate of 37.4 km. The estimated detection probability was 0.54129 (%CV = 8.07; df = 434.00; 95 % CI = 0.461 99–0.634 20). The maximum observed perpendicular distance to active

burrow entrances was 10 m. The total area surveyed by transects was 23.38 ha. The effective strip width (ESW) was 5.4 m (CV = 8.1 %, 95 % CI = 4.6–6.3 m). The density of active burrow openings was 34.53 per hectare (SE = 3.44, %CV = 9.95).

The 2023 survey yielded 162 active burrow openings observations, with an encounter rate of 8.67 km. The detection probability was estimated at 0.24527 (%CV = 12.95 %; df = 158; 95 % CI = 0.190 11–0.316 45) and 29 transects registered no observations. The maximum observed perpendicular distance to active burrow entrances was 7.5 m and the total area surveyed was 28.03 ha. The effective strip width (ESW) was 1.8 m (CV = 13.0 %, 95 % CI = 1.4–2.3 m). The density of active burrow openings was 23.55 per hectare (SE = 5.45, %CV = 23.14).

Density data for each location and year are presented in Tab. 1.

The effective strip width (ESW) estimated in 2023 was substantially lower than in 2022, indicating a steeper decline in detectability with increasing distance from transect lines. This difference may reflect surveys being conducted earlier in the vegetation season in 2023, when herbaceous cover is generally denser, potentially reducing visibility of burrow entrances at greater distances relative to late summer conditions in 2022. However, vegetation structure was not quantified directly, and this interpretation should be regarded as contextual rather than causal.

Interannual differences in encounter rate and detection probability further reflects differences in sampling strategies. In 2022, surveys intentionally focused within known colonies, where burrow entrances are spatially clustered (Gedeon et al., 2022; Kachamakova et al., 2022), resulting in high encounter rates. In contrast, the 2023 surveys employed random transects distributed across a broader landscape, including areas with low or no burrow presence, which led to reduced encounter rates and a higher proportion of transects without detections. These design differences are reflected in lower detection probability estimated for 2023.

Density estimates mirrored these methodological contrasts. The systematic, colony-focused design used in 2022 yielded more precise density estimates, but it primarily characterizes conditions within occupied colonies. In contrast, the random transect design applied in 2023 provides density estimates that are more representative of the broader habitat matrix, albeit with increased variability. The higher coefficient of variation observed in 2023 indicates greater uncertainty, consistent with the inclusion of sparsely occupied habitat and a lower number of detections contributing to detection-function estimation. When analysed separately by location and year, observations were often too few, resulting in wide confidence intervals and imprecise density estimates. These results are reported for completeness but should be interpreted cautiously due to low sample sizes in some cases. To enable meaningful interpretation and improve data reliability in future surveys, it is recommended to increase survey effort.

Distance sampling offers a robust statistical framework for estimating densities of EGS active burrow openings by explicitly modelling detectability as a function of distance (Buckland et al., 2001). Empirical studies have shown that counts of active burrow entrances are positively correlated with individual abundance in this species (Katona et al., 2002; Stoeva et al., 2016; Gedeon et al., 2017), supporting the

**Table 1** – Location and year-specific density estimate of active burrow openings derived from conventional distance sampling and the mapped suitable habitat area (SE - Standard error; %CV - Coefficient of variation; 95% CI - Confidence interval).

| Year | Location             | Density (active burrow openings/ha)        | SE     | %CV   | 95% CI          | Suitable habitat area (ha) |
|------|----------------------|--------------------------------------------|--------|-------|-----------------|----------------------------|
| 2022 | Sânpetru German (SG) | 34.184                                     | 4.5697 | 13.37 | 26.283 - 44.462 |                            |
|      | Felnac (FC)          | 32.208                                     | 3.5814 | 11.12 | 25.795 - 40.214 |                            |
|      | Zădăreni (ZD)        | 27.927                                     | 6.4622 | 23.14 | 17.574 - 44.378 |                            |
|      | Igrîș (IG)           | 31.827                                     | 18.319 | 57.69 | 9.6905 - 104.53 | 52                         |
|      | Secusigiu (SC)       | 43.591                                     | 16.498 | 37.35 | 19.834 - 95.805 | 229                        |
| 2023 | Munar (MN)           | Not estimated, not enough observation data |        |       |                 | 256                        |
|      | Felnac (FC)          | 19.966                                     | 5.8869 | 29.48 | 11.192 - 35.620 | 267                        |
|      | Zădăreni (ZD)        | 12.916                                     | 5.9697 | 46.22 | 4.8728 - 34.236 | 60.2                       |



use of burrow-based distance sampling as a reliable approach for population assessment. 

## References

- Buckland S.T., Anderson D.R., Burnham K.P., Laake J.L., Borchers D.L., & Thomas L., 2001. *Introduction to Distance Sampling: Estimating Abundance of Biological Populations*. Oxford University Press, Oxford. 10.1093/OSO/9780198506492.001.0001
- Burnham K.P. & Anderson D.R., 2004. Multimodel Inference: Understanding AIC and BIC in Model Selection. *Sociological Methods & Research*, 33(2): 261–304. <https://doi.org/10.1177/0049124104268644>
- Čosić N., Čirović D., Fülöp T., Gedeon C., Hapl E., Hoffmann I.E., Kepel A., Koshev Y., Matějů J., Nikolić Lugonja T., Rammou L.-D., Rusin M., Váczi O., & Hegyeli Z., 2024. *Spermophilus citellus*. *The IUCN Red List of Threatened Species* 2024: e.T20472A221789466. <https://dx.doi.org/10.2305/IUCN.UK.2024-2.RLTS.T20472A221789466.en> (Accessed on 24 February 2025).
- E.E.A., 2025. *Environment Agency Reportnet. Article 17 web tool on biogeographical assessments of conservation status of species and habitats under Article 17 of the Habitats Directive*. <https://nature-art17.eionet.europa.eu/article17/> (Accessed on 24 February 2025).
- Gedeon C.I., Árvai M., Szatmári G., Brevik E.C., Takáts T., Kovács Z.A., & Mészáros J., 2022. Identification and Counting of European Souslik Burrows from UAV Images by Pixel-Based Image Analysis and Random Forest Classification. *Remote Sens.*, 14(9). <https://doi.org/10.3390/rs14092025>
- Gedeon C.I., Hofmann I.E., Váczi O., Knauer F., Slimen H.B., Stepanovic M., Lehoczki É., Laborczí A., & Suchentrunk F., 2017. The role of landscape history in determining allelic richness of European ground squirrels (*Spermophilus citellus*) in Central Europe. *Hystrix It. J. Mamm.*, 28(2): 231–239. 10.4404/hystrix-28-2-11823
- Hegyeli Z., Nagy A., Daroczi S.J., Kecskes A., & Latkova H., 2012. Current distribution and status of the European ground squirrel in Romania. In: Kepel A. & Kończak J. (Eds.), *IV European Ground Squirrel Meeting, Kamień Śląski, Poland*. Polish Society for Nature Conservation “Salamandra”.
- Ionescu O., Ionescu G., Jurj R., Cazacu C., Adamescu M., Cotovelea A., Pașca C., Popa M., Mirea I., Sîrbu G., Chiriac S., Pop M., Atilla Ş., & Deju R., 2013. *Ghid sintetic de monitorizare pentru specile de mamifere de interes comunitar din România*. Editura Silvică, Voluntari.
- Jacquier M., Vandel J.M., Léger F., Duhayer J., Pardonnet S., Say L., Devillard S., Ruette S., 2021. Breaking down population density into different components to better understand its spatial variation. *BMC Ecol. Evol.*, 21(1): 82. 10.1186/S12862-021-01809-6
- Kachamakova M., Koshev Y., Rammou D.L., & Spasov S., 2022. Rise and fall: Results of a multidisciplinary study and 5-year long monitoring of conservation translocation of the European ground squirrel. *Biodivers. Data J.*, 10: e83321. 10.3897/BDJ.10.E83321
- Katona K., Váczi O., & Altbäcker V., 2002. Topographic distribution and daily activity of the European ground squirrel population in Bugacpuszta, Hungary. *Acta Theriol.*, 47(1): 45–54. 10.1007/BF03193565
- Koshev S.Y., 2008. Distribution and status of the European ground squirrel (*Spermophilus citellus*) in Bulgaria. *Lynx (Praha)*, n. s., 39(2): 251–261.
- Kryštufek B., 1999. *Spermophilus citellus* (Linnaeus, 1766). In: Mitchell-Jones A.J. (Ed.). *The atlas of European mammals*. Academic Press, London, pp. 190–191.
- Kryštufek B. & Vohralík V., 2012. Taxonomic revision of the Palaearctic rodents (Rodentia). Sciuridae: Xerinae 1 (*Eutamias* and *Spermophilus*). *Lynx, n.s. (Praha)*, 43(1-2): 17–111.
- Luikart G., Ryman N., Tallmon D.A., Schwartz M.K., & Allendorf F.W., 2010. Estimation of census and effective population sizes: The increasing usefulness of DNA-based approaches. *Conservation Genetics*, 11(2): 355–373. 10.1007/S10592-010-0050-7
- MMAP, 2016. *Planul de management al Parcului Natural Lunca Mureșului* [Management Plan of the Lunca Mureșului Natural Park] (Anexă din 30 iunie 2016). Monitorul Oficial, Partea I, nr. 836bis. București.
- Murariu D., 2005. Mammalia (Mamifere). In: Botnariuc N. & Tatole V. (Eds.) *Cartea roșie a vertebratelor din România*. Ed. Muzeul Național de Istorie Naturală “Gr. Antipa”, București, pp. 11–84.
- Rammou D.-L., Kavrouidakis D., & Youlatos D., 2021. Distribution, Population Size, and Habitat Characteristics of the Endangered European Ground Squirrel (*Spermophilus citellus*, Rodentia, Mammalia) in Its Southernmost Range. *Sustainability*, 13: 8411. 10.3390/su13158411
- Ramos-Lara N., Koprowski J.L., Kryštufek B., & Hoffmann I., 2014. *Spermophilus citellus* (Rodentia: Sciuridae). *Mammalian Species*, 46(913): 71–87. 10.1644/913.1
- Ružić A., 1978. *Citellus citellus* (Linnaeus, 1766) - der Oder das Europäische Ziesel. In: Niethammer J. & Krapp F. (Eds.) *Handbuch der Säugetiere Europas*. Bd. 1, Nagetiere I. Akademische Verlagsgesellschaft, Leipzig, Germany, pp. 123–144.
- Stoeva E., Ivanov I., Stoev I., Yankov L., Mechev A., & Koshev Y., 2016. Successful reinforcement of the European souslik by Green Balkans NGO in “Sinite Kamani” Nature Park, Bulgaria. *Annuaire de l'Université de Sofia “St. Kliment Ohridski” Faculte de Biologie*, 153–165.
- Thomas L., Buckland S.T., Rexstad E.A., Laake J.L., Strindberg S., Hedley S.L., Bishop J.R.B., Marques T.A., & Burnham K.P., 2010. Distance software: design and analysis of distance sampling surveys for estimating population size. *Journal of Applied Ecology*, 47: 5–14. 10.1111/j.1365-2664.2009.01737.x
- Thompson W., 2004. *Sampling rare or elusive species: Concepts, designs, and techniques for estimating population parameters*. Island Press, Washington, D.C.

Associate Editor: G. Amori