

## POST CONSTRUCTION MONITORING OF WIND FARMS: FIRST RECORDS OF DIRECT IMPACT ON BATS IN ITALY

VINCENZO FERRI<sup>1\*</sup>, OSVALDO LOCASCIULLI<sup>2</sup>, CHRISTIANA SOCCINI<sup>1</sup>,  
ELIA FORLIZZI<sup>2</sup>

<sup>1</sup>Centro Studi Arcadia, Via Valverde 4, 01016 Tarquinia, Viterbo, Italy

\*Corresponding author, E-mail: vincenf@tin.it

<sup>2</sup>Riserva Naturale Regionale "Lago di Penne", Corso dei Vestini 4,  
65017 Penne, Pescara, Italy

Received 6 October 2010; accepted 20 April 2011

**ABSTRACT** - Long-term post-construction monitoring of the impact of wind farms on fauna started in 2008 in Abruzzo (central Italy). This study provides the first evidence of bat fatalities at wind turbines in Italy. Six carcasses of *Hypsugo savii* and one carcass of *Pipistrellus pipistrellus* were recorded in 2009 during 36 field sessions, involving 4 researchers at two wind farms, for a total of 46 (21 and 25) wind turbines.

**Key words:** wind farms, monitoring, impact, bats, Italy

**Riassunto - Monitoraggi post-operam di impianti eolici: prima segnalazione di impatto diretto su pipistrelli in Italia.** La possibilità di attuare a partire dal 2008 un monitoraggio post-operam della fauna di interesse conservazionistico presso impianti eolici in Abruzzo ha permesso di evidenziare i primi casi certi di fatalità di pipistrelli in Italia. Durante 36 sessioni di campo del 2009, da parte di 4 operatori, presso due impianti eolici industriali, composti il primo di 21 ed il secondo di 25 turbine, sono state rilevate 6 carcasse di *Hypsugo savii* e una di *Pipistrellus pipistrellus*.

**Parole chiave:** monitoraggio post-operam, impianti eolici, impatto, pipistrelli, Italia

DOI: 10.4404/Hystrix-22.1-4525

### INTRODUCTION

The uncontrolled increase of greenhouse gases and the increasingly tangible negative effects of global warming are resulting in an ever-stronger demand for more sources of renewable energy (Solomon et al. 2007). In little more than 20 years, industrial wind farms have increased exponentially, both in Europe and worldwide. Hazards and drawbacks of wind energy devel-

opment for the environment and especially for wildlife are under discussion. However, it is clear that birds and bats are the most threatened groups due to collisions and close-contact injuries. Since the end of the 90's several studies on flying wildlife, from Europe and North America, have shown some occurrence of bat mortality at wind farms, because of the direct impact with the spinning wind turbine blades (Lekuona 2001; Erickson et al. 2003; Arnett

2005); in some cases the number of bats involved per year has been higher than that of birds, overall. Bat mortality data at wind farms are updated every year both by EUROBATS Intersessional Working Group (AAVV 2010) and by Dürr and Bach (2004) and Dürr (2007).

The majority of kills involved potential tree bat species (i.e. species that roost within the foliage or in tree cavities) and the impact peak was found in late summer and autumn (Kerns et al. 2004). Collision and mortality peaks coincide with their migration and mating periods, but European studies showed a significant impact also on non-migratory species, such as *Pipistrellus* spp. (Rodrigues et al. 2008). A recent comprehensive world-wide analysis on the different types of mortalities within industrial wind farms (Cryan and Barclay 2009) confirmed that the most affected species are those linked to trees and which often migrate across latitudes. However, the reason why individuals of those and other non-migratory species end up crashing against the turbines is still unclear. Determining the causes of bat fatalities at turbines is important in order to develop practical solutions to this serious problem for bat conservation. Here we present the preliminary results of post-construction bat fatality searches that are part of a long-term monitoring (2009-2012) strategy at wind farms in Abruzzo (Central Italy).

## MATERIALS AND METHODS

The study area is located near the Fucino Valley and the Sirente-Velino Natural Regional Park, along the southern slopes of the Sirente Massif, with an altitudinal range

of 900-1150 m a.s.l. (WGS 84, F33 coordinates: top left - X 381.985, Y 4.663.406; bottom right - X 397.520, Y 4.653.612). The post-construction monitoring considered the wind farm of Cerchio-Collarmele (WF1) that began operation in March 2009 and consists of 21 2.0 MW turbines (Vestas V80) and 3 meteorological towers. Rotor diameter of each turbine is 80 m and spans over an area of 5072 m<sup>2</sup> and the tubular steel towers are 78 m high. Therefore, each turbine reaches a maximum height, at the tip of the blades, of 117 m from ground level. Non periodical surveys were also carried within the nearby Cocullo Wind Farm (WF2), 15 km away. WF2 was built in 2005 and consists of 38 Gamesa 850 kW turbines. This site is located along a mountain ridge between 1200 m and 1600 m a.s.l.

At both sites, the natural vegetation where wind farms are located consists of high hill - low mountain scrubland and hemi-cryptophytic pasture patches characterised by *Brachypodium rupestre*. In the sites with cattle present there are nitrophilous phyto-coenoses and sparse bushes (e.g. *Rosa canina*, *Prunus spinosa*, *Rubus ulmi-folius*, *Crataegus monogyna*).

The surrounding areas of 46 turbines (21 at WF1 and 25 at WF2) were searched for bat carcasses every three days from March 15th to October 31st 2009, with 32 field sessions in WF1 and 4 field sessions in WF2. Permanent square plots, 120 m per side and centred on the turbine were set up. The searcher walking at a rate of approximately 45-60 m/min along each transect of 120 m, searching both sides up to 3 m for bat carcasses (Johnson et al., 2000, 2003). Subsequently, both transect width and search speed were based on visibility within the various habitat types and vegetation stages. On average, approximately 30 to 60 minutes were spent searching each turbine, depending on the habitat type. For each carcass found, recorded data included species, sex and age class (when discernable), date and time of collection, location

(WGS 84 F33 coordinates), habitat type, condition, and any further comment which may help estimating time and cause of death.

## RESULTS AND DISCUSSION

During field sessions seven bat carcasses were found: 1 *Pipistrellus pipistrellus* and 6 *Hypsugo savii*. Dead bats were all found from 3 August to 16 September (Tab. 1); 3 carcasses were found beneath one of the WF1 turbines and 4 carcasses beneath four of the WF2 turbines. Three of the collected bat carcasses were fresh and intact: necropsy showed abundant haemorrhages within both the abdominal and the thoracic cavities, referable to the syndrome described as “baro-trauma” (Baerwald et al. 2008). Three other carcasses were scavenged by wasps and ants and one had been dismembered. The distances from turbines where the dead bats were found ranged from 0.7

m to 14.4 m and the average was 14.2 m. All of the bats were found during fair weather conditions.

In this early phase of post-construction monitoring at our study sites in Italy, we are not yet able to accurately estimate fatality rates and thus make comparisons with other European studies, especially in cases where carcasses of different species are encountered. However, we have documented for the first time the fatality of both *Pipistrellus pipistrellus* and *Hypsugo savii* at wind turbines in Italy.

Both of these species have been found dead beneath turbines in other European countries, also mostly during late summer and fall. *P. pipistrellus* and *H. savii* are not known to migrate in Italy and presumably non-migratory populations also show susceptibility to turbines as in other parts of Europe (e.g., Germany; Dürr and Bach 2004).

Monitoring at our study sites will con-

Table 1 - Bat carcasses found in 2009 during post-construction monitoring of Abruzzo’ wind farms WF1 “Cerchio-Collarmele” and WF2 “Cocullo”; M: male; F: female; n.d.: not determined.

Species	Wind Farms	Date	Carcass condition	Distance from turbine (m)	Sex	Forearm length (mm)	Weight (g)
<i>P. pipistrellus</i>	WF2 Cocullo	03.09.09	partially scavenged by wasps / dried	1.40	n.d.	27.34	1.1
<i>H. savii</i>	cluster 1 WF1	03.08.09	intact / fresh	6.50	M	33.44	4.9
<i>H. savii</i>	WF2 Cocullo	09.08.09	partially scavenged by ants / dried	14.40	M	33.04	1.8
<i>H. savii</i>	cluster 1 WF1	27.08.09	intact / fresh	2.40	F	34.22	5.5
<i>H. savii</i>	cluster 1 WF1	11.08.09	intact / fresh	8.00	n.d.	n.d.	1.0
<i>H. savii</i>	WF2 Cocullo	03.09.09	rests of carcass scavenged by ants	11.00	n.d.	n.d.	1.2
<i>H. savii</i>	WF2 Cocullo	16.09.09	disjointed	8.20	n.d.	n.d.	0.8

tinue for two more years and final results, corrected for searcher efficiency, scavenger removal, and search-interval bias (Huso 2010) could provide a significant contribution to the understanding of the situation with bats and wind energy in Italy. The owners of WF1 have already been involved with this problem and we have proposed a mitigation activity to reduce bat mortality, as realized in some wind farms in Canada (Baerwald et al., 2009), Pennsylvania (Arnett et al. 2009) and Germany, by changing the cut-speed (the minimum wind speed when the rotors start spinning) from a “normal” limit (3.5 - 4.0 m/sec) to 5.5 m/sec.

## ACKNOWLEDGMENTS

The authors are grateful to the persons in charge and the technicians of the wind farm that made it possible to operate and carry out the research. The Sirente -Velino Natural Regional Park authorised the studies (Prot. 2666) as well as the Ministry of the Environment and the ISPRA (authorisation received on 25.06.2007 (DPN-2007-0017838). Paul Cryan and Marie-Jo Dubourg-Savage improved the paper with corrections and useful suggestions.

## REFERENCES

AAVV 2010. Report of the IWG on Wind Turbines and Bat Populations. Doc.EUROBATS.StC4-AC15.22.Rev.1. 4<sup>th</sup> Meeting of the Standing Committee. 15<sup>th</sup> Meeting of the Advisory Committee. Bonn, Germany. Available from: [http://www.eurobats.org/documents/pdf/AC15\\_StC4/Doc\\_StC4\\_AC15\\_22\\_Rev1\\_ReportIWG\\_WindTurbines\\_incl\\_annexes.pdf](http://www.eurobats.org/documents/pdf/AC15_StC4/Doc_StC4_AC15_22_Rev1_ReportIWG_WindTurbines_incl_annexes.pdf). [20 September 2010].

Arnett E.B. (Ed.) 2005. Relationships between bats and wind turbines in Penn-

sylvania and West Virginia: an assessment of bat fatality search protocols, patterns of fatality, and behavioral interactions with wind turbines. A final report submitted to the Bats and Wind Energy Cooperative. Bat Conservation International, Austin, Texas, USA. Available from: <http://www.batcon.org/wind/BWEC2004finalreport.pdf>. [1 May 2007].

Arnett E.B., Schirmacher M., Huso M.M.P., Hayes J.P. 2009. Effectiveness of Changing Wind Turbine Cut-in Speed to Reduce Bat Fatalities at Wind Facilities. Annual Report Prepared for the Bats and Wind Energy Cooperative and the Pennsylvania Game Commission.

Baerwald E.F., D’amours G.H., Klug B.J., Barclay R.M.R. 2008. Barotrauma is a significant cause of bat fatalities at wind turbines. *Curr. Biol.* 18: R695–696.

Baerwald E.F., Edworthy J., Holder M., Barclay R.M.R. 2009. A large-scale mitigation experiment to reduce bat fatalities at wind energy facilities. *J. Wildl. Manage.* 73: 1077–1081.

Crawford R.L., Baker W.W. 1981. Bats killed at a north Florida television tower: a 25-year record. *J. Mammal.* 62: 651–652.

Cryan P.M., Barclay R.M.R. 2009. Causes of bat fatalities at wind turbines: hypotheses and predictions. *J. Mammal.* 90(6): 1330–1340

Dürr, T., Bach L. 2004. Fledermäuse als Schlagopfer von Windenergieanlagen - Stand der Erfahrungen mit Einblick in die bundesweite Fundkartei. *Bremer Beiträge für Naturkunde und Naturschutz* 7: 253–264.

Dürr T. 2007. Die bundesweite Kartei zur Dokumentation von Fledermausverlusten an Windenergieanlagen - ein Rückblick auf 5 Jahre Datenerfassung. *Nyctalus (N.F.)* 12(2/3): 108–114.

Erickson W.P., Gritski B., Kronner K. 2003. Nine Canyon Wind Power Pro-

- ject Avian and Bat Monitoring Report, September 2002 - August 2003. Technical report submitted to Energy Northwest and the Nine Canyon Technical Advisory Committee.
- Hall L.S., Richards G.C. 1972. Notes on *Tadarida australis* (Chiroptera: Molossidae). *Aust. Mammal.* 1:46.
- Huso M.M.P. 2010. An estimator of wildlife fatality from observed carcasses. *Environmetrics*. Published online in Wiley InterScience. [www.interscience.wiley.com](http://www.interscience.wiley.com) doi: 10.1002/env.1052.
- Kerns J, Kerlinger P. 2004. A study of bird and bat collision fatalities at the Mountaineer Wind Energy Center, Tucker County, West Virginia. Annual report for 2003. Curry and Kerlinger, LLC, McLean, Virginia, USA. Available from: <http://www.batcon.org/windliterature>. [1 September 2007].
- Johnson G.D., Erickson W.P., Strickland M.D., Shepherd M.F., Shepherd D.A. 2000. Avian Monitoring Studies at the Buffalo Ridge Wind Resource Area, Minnesota: Results of a 4-year study. Technical Report prepared by WEST, Inc. for Northern States Power Co., Minneapolis, MN. 212 pp.
- Johnson G.D., Erickson W., White J., McKinney R. 2003. Avian and bat mortality during the first year of operation at the Klondike. Phase I Wind Project, Sherman County, Oregon. March 2003. Tech. Rept. prepared for West Inc., Cheyenne 17 pp.
- Lekuona J.M. 2001. Uso del espacio por la avifauna y control de la mortalidad de aves y murciélagos en los parques eólicos de Navarra durante un ciclo anual. Dirección General de Medio Ambiente, Gobierno de Navarra.
- Osborn R.G., Higgins K.F., Dieter C.D., Usgaard R.E. 1996. Bat collisions with wind turbines in Southwestern Minnesota. *Bat Research News* 37: 105-108.
- Rodrigues L., Bach L., Dubourg-Savage M.-J., Goodwin J., Harbusch C., 2008. Guidelines for consideration of bats in wind farm project, EUROBATS Publication Series No. 3 (English version). UNEP/EUROBATS Secretariat, Bonn, Germany, 51 pp.
- Solomon S., Qin D., Manning M., Chen Z., Marquis M., Averyt K.B., Tignor M., Miller H.L. (Eds.) 2007. IPCC. Climate Change 2007: The Physical Science Basis. Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA, 996 pp.
- Strickland M.D., Erickson W.P., Johnson G., Young D., Good R. 2001. Risk reduction avian studies at the Foote Creek Rim Wind Plant in Wyoming. Proceedings of the National Avian-Wind Power Planning Meeting IV. National Wind Coordinating Committee, c/o RESOLVE, Inc. Washington, D.C.
- Van Gelder R.G. 1956. Echo-location failure in migratory bats. *Transactions of the Kansas Academy of Science* 59:220-222.