



Available online at:

<http://www.italian-journal-of-mammalogy.it/article/view/12421/pdf>

doi:10.4404/hystrix-28.2-12421

Short Note

## From accidental citizen-science observations to genetic confirmation: how to spot new hidden invaders

Emiliano MORI<sup>1,\*</sup>, Alberto BAERI<sup>1</sup>, Andrea SFORZI<sup>2</sup>, Alessandro VITALE<sup>3</sup>, Andrea GALIMBERTI<sup>4</sup><sup>1</sup>Dipartimento di Scienze della Vita, Università di Siena, Via P.A. Mattioli, 4 - 53100 Siena, Italy<sup>2</sup>Museo di Storia Naturale della Maremma, Strada Corsini, 5 – 58100 Grosseto, Italy<sup>3</sup>Department of Prevention, Food Hygiene and Nutrition Service ASUR Marche A.V.3 Macerata, Via Belvedere Sanzio 1, 62100 Macerata.<sup>4</sup>ZooPlantLab, Department of Biotechnology and Biosciences, University of Milano-Bicocca, P.zza della Scienza 2, 20126-I Milano, Italy

### Keywords:

*Hystrix cristata*  
*Hystrix indica*  
 molecular identification  
 DNA barcoding

### Article history:

Received: 1 March 2017

Accepted: 23 June 2017

### Acknowledgements

Authors thank two anonymous reviewers for their useful suggestions on an early draft

### Abstract

The crested porcupine *Hystrix cristata* is the largest rodent in Italy, where it is strictly protected according to national and international laws. The species is almost exclusively nocturnal and quite elusive: its presence may be yet recorded through the detection of quills lost on the ground. In the last 40 years, the crested porcupine showed a remarkable range expansion in Italy. In 2013, in the framework of the [www.naturaesocialmapping.it](http://www.naturaesocialmapping.it) recording site, a web page devoted to *H. cristata* was set up. The aim of the project was to give people the opportunity to contribute to a continuous updating of the distributional map of this rodent. Lost quills were also collected by citizen-observers, as being easily available sources of DNA. This web page allowed us to detect the first evidence of the presence of another species of porcupine free-ranging in Italy (photos and quills): the Indian porcupine *H. indica*. Species identification was carried out through DNA barcoding analysis, since it cannot be performed only through quill morphology. We increased the international genetic database with sequences of both species, so to allow a rapid identification of samples through molecular analyses. Both species were introduced to Italy by humans. Our findings also support the relevance of citizen science contribution in the process of updating species' distribution range and the potential of spotting new invasive species.

The systematic status and field identification of many rodent taxa are still controversial and debated, especially when species belonging to the same genus share similar morphological features and geographical range (Colangelo et al., 2009; Castiglia et al., 2016; Mazzamuto et al., 2016; Wauters et al., 2017). In most cases, samples (e.g., hairs, bones and faeces) found during field surveys could not provide sufficient information to achieve unequivocal identification. In this context, advances in molecular DNA-based tools represent a basic element to improve knowledge in taxonomy (Corti et al., 2005; Colangelo et al., 2009; Abiadh et al., 2010; Padial et al., 2010; Ermakov et al., 2015), that complement monitoring activities. Many investigations in the field of bioinvasions are currently conducted under an integrative vision, where variability of molecular markers is used in combination with morphological traits, geographical origin and other sources of information (Gotzek et al., 2012; Pisanu et al., 2013; Mazzamuto et al., 2016). In the last years, DNA barcoding triggered a real revolution inside species identification, with applications ranging from pure taxonomy to cases of conservation or forensics concern (Hebert et al., 2003) This identification approach has been successfully applied on a vast plethora of case studies involving mammals (Galimberti et al., 2015). Publicly accessible archives including reference DNA barcoding sequences (targeting especially at region of the mitochondrial *COI*) are growing, thus offering the possibility to confirm doubtful field observations, even using non-invasive sampling approaches.

Porcupines are nocturnal, elusive rodents and most sightings are related to observations of (i) quills lost on the ground, (ii) dens and (iii) diggings on the ground to get food (Mori et al., 2013). The subgenus *Hystrix* includes three genetically related species of porcupines, distributed throughout Africa, Southern Europe and Asia with areas of almost

complete allopatry (Mohr, 1965; Walid, 2011; Trucchi et al., 2016). The crested porcupine *H. cristata* L., 1758 shows a discontinuous geographical range. It is distributed in a portion of Southern Europe (peninsular Italy and Sicily, introduced in Sardinia), Northern Africa (from Morocco to Libya) and sub-Saharan Africa (from Senegal to Tanzania). In Central and Southern Tanzania, the range of *H. cristata* partly overlaps with the range of the Cape porcupine *H. africae australis* Peters, 1852 (Skinner and Smithers, 1990; Barthelmess, 2006). The Indian porcupine *H. indica* Kerr, 1792 ranges from Turkey and Israel to Western India and Nepal (Mohr, 1965). These species are morphologically very similar, and very difficult to set apart when syntopic. Main diagnostic features are summarized in Tab. 1 (Corbet and Jones, 1956; Barthelmess, 2006; Aulagnier et al., 2010).

The range expansion of the crested porcupine in Italy has been recorded through a citizen-science approach, still ongoing. In 2013, in the framework of the [www.naturaesocialmapping.it](http://www.naturaesocialmapping.it) recording site, a web page devoted to *H. cristata* was set up. The aim of the project was to give people the opportunity to contribute to a continuous updating of the distributional map of this rodent. This approach has been proven to be very successful, as the species is unmistakable, with just a 0.03% of identification error. In September 2016, an individual showing the outer morphology of an Indian porcupine, i.e. a black crest, was accidentally recorded in the surroundings of a den in Abbadia di Fiastra (Province of Macerata, Central Italy), representing the first formal observation of this species in the wild outside its natural range. Quills were collected and genetically analyzed through DNA barcoding for the identification of the species.

We therefore produced the first reference DNA barcoding records for *H. cristata*. DNA barcoding sequences of *Hystrix cristata* were not included in the international GenBank (<https://www.ncbi.nlm.nih.gov>) and the Barcode of Life Data System (<http://www.boldsystems.org/>),

\*Corresponding author

Email address: moriemiliano@tiscali.it (Emiliano MORI)

**Table 1** – Main diagnostic features in the subgenus *Hystrix*. MAR, midline along the rump; NB, nasal bones; ONL, occipito-nasal length; P, premaxilla; PW, premaxilla width; NBW, nasal bone width (from Corbet and Jones, 1965; Barthelmeß, 2006; Aulagnier et al., 2010).

Species	Colour of the crest	Colour of MAR	NB (% of ONL)	P (% of PW/NBW)
<i>Hystrix cristata</i>	predominantly white	black or mottled	long (58-68)	intermediate (< 36)
<i>Hystrix indica</i>	predominantly brown	white	short (45-50)	wide (> 44)
<i>Hystrix africaeaustralis</i>	predominantly white	white	intermediate (51-58)	narrow (< 23)

whereas only four *COI* sequences of *H. indica* were available (databases last accessed on March 2017). Thus, a robust reference DNA barcoding dataset was created for both *Hystrix* species to achieve a reliable identification of the samples found in Central Italy (hereafter referred to as “Macerata”). A total of eight samples belonging to *H. cristata*, *H. indica* and the quills from the individual found at Abbadia di Fiastra, were collected (Tab. 2). Reference samples belonged to individuals of *Hystrix cristata* collected in Italy (captures or road-killed), and to individuals of *Hystrix indica* from Turkey and Israel (road-killed). To maximize the chance of observing intraspecific geographic variation, conspecifics were sampled from distant sites (Table 2). Samples were vouchered and then stored in 99% ethanol at -20°C.

Total DNA was extracted from 25 mg of tissue or from the basal portion (4 mm) of three quills by using the DNeasy Blood & Tissue Kit (Qiagen, Milan, Italy) following manufacturer’s instructions with minor modifications in the case of quills. In particular, lysis was performed overnight and final DNA was eluted in 40 µl of dehyonized sterile water warmed up to 65°C. Purified DNA concentration of each sample was estimated fluorometrically with a NanoDrop™ 1000 Spectrophotometer (Thermo Scientific, USA). DNA barcoding characterization of samples was conducted by comparing nucleotide sequence differences at the standard barcode region for metazoans (i.e. 648 bp at the 5’ end of the mitochondrial *COI*: Hebert et al. (2003)) following the same procedures described in Mazzamuto et al., 2016. Sequence data were submitted to the European Bioinformatics Institute of the European Molecular Biology Laboratory (EMBL-EBI). In addition to the samples analyzed in this study, we considered the four DNA barcoding sequences of *H. indica* publicly available in GenBank and an outgroup *COI* sequence of *Atherurus africanus* (a.n. KJ192744). Sequences were aligned using the ClustalW option implemented in Bioedit (Hall, 1999) with default options. The taxonomic status of the Macerata quills sample was also tested by comparing the *COI* sequence with reference DNA barcoding stored in the Barcode of Life Database using the Identification Engine tool (IDS) ([http://www.boldsystems.org/index.php/IDS\\_OpenIdEngine](http://www.boldsystems.org/index.php/IDS_OpenIdEngine); Species Level Barcode Records database), which returns unique species assignments based on 99% sequence similarity of the barcode sequence. Considering the comprehensive dataset, average K2P genetic sequence divergences (and relative standard errors, SE) between and within distinct lineages were calculated and a NJ reconstruction was performed using MEGA 6 with the same settings described in Galimberti et al., 2012.

DNA extracted from the ethanol preserved tissues and quills was of high quality (ratios of absorbance,  $A_{260/280}$  and  $A_{260/230} \sim 1.80$  and  $>1.90$ , respectively) and provided good yields ( $> 30$  ng/µl). Amplification with the selected primer pairs was successful and resulting DNA concentration of purified amplicons was  $>50$  ng/µl. High quality sequences showing a strong chromatogram signal along the entire read were obtained for all the individuals sampled for the present study. Due to different lengths of GenBank sequences, we trimmed the alignment to the same final size of 645 bp. The *COI* DNA barcoding dataset contained sequences with no insertion/deletions (indels), stop codons or biased by NUMT (*sensu* Bensasson et al. (2001)) interference. The alignment was characterized by 64 variable positions, of which 62 were parsimony-informative. Despite different sampling localities, all samples of *H. cristata* shared the same haplotype (HAP1), whereas, four haplotypes were found for *H. indica*, one of which shared by three Genbank entries (HAP5) and the other three exclusive to the remaining accessions (i.e., HAP6 from GenBank and HAP3 and HAP4 from the two samples analyzed in this study, including the sample of Macerata). The Macerata *Hystrix* sample showed an exclusive *COI* hap-


lotype (HAP4). Sequences accessions and haplotypes are provided in Tab. 2. When *COI* sequence obtained from the Macerata sample was used as query in GenBank-NCBI and BOLD-IDS

tool, both systems returned a reliable match (i.e., similarity %  $> 99\%$ ) with reference DNA barcoding sequences of *H. indica*. The analysis of the K2P distance matrix (Table S1) revealed a moderate intraspecific divergence among lineages at the *COI* locus for *H. indica* (mean K2P distance  $\pm$  standard error:  $1.25\% \pm 0.31\%$ , range: 0 - 2.06%) and a large interspecific divergence between *H. cristata* and *H. indica* (mean K2P distance  $\pm$  standard error:  $9.38\% \pm 1.26\%$ , range: 9.06 - 9.80%). Despite the investigation of the other five *Hystrix* species should be recommended to assess the actual genetic distances, the existence of a clear barcoding gap (Wiemers and Fiedlers, 2007) may confirm the efficacy of DNA barcoding in unambiguously identifying *Hystrix* species.

The Macerata *Hystrix* sample had HAP2 and HAP6 haplotypes belonging to *H. indica* as its nearest neighbors (K2P distance  $\pm$  standard error:  $0.16\% \pm 0.15\%$ ), whereas a much higher genetic divergence occurred between the Macerata sample and *H. cristata* haplotypes (K2P distance  $\pm$  standard error:  $8.88\% \pm 1.23\%$ ).

The NJ reconstruction (Fig. 1) better summarized the pattern of molecular variability and unequivocally supported the assignment of the Macerata quill sample to the species *H. indica*.

In our work, we confirmed the efficacy of a standard molecular method, which allows for a rapid identification of porcupine species where they occur syntopically. The high molecular divergence at the barcode locus *COI* between the two *Hystrix* species (i.e.  $> 8\%$  K2P) makes DNA barcoding approach reliable to assess the occurrence of the invasive taxon also starting from samples retrieved from non-invasive approaches. Such a molecular divergence is comparable to values obtained in previous studies targeting at *Cyt b* (K2P distance 10.38%) and *D-loop* (K2P 8.22%–9.21%) mitochondrial regions (Trucchi et al., 2008, 2009). However, such values are biased by the analysis of a single *H. indica* sample (compared to the 7 used in our DNA barcoding approach) and further investigations are needed to better support genetic distances at these markers. In addition, the high number of nucleotide differences at the barcode locus between *H. cristata* and *H. indica* permits the development of fast (and sequencing free) detection



**Figure 1** – Neighbour joining tree based on *COI* sequences of *Hystrix cristata* and *H. indica*. For each samples, voucher or GenBank accession number is provided (further details can be retrieved from Tab. 2). Bootstrap support (1000 replicates) values  $> 90\%$  are indicated above the nodes. The filled circle shows the Macerata quill sample. A colour version of this figure can be viewed in the online version of this article.

**Table 2** – Reference samples of *H. cristata* and *H. indica*, with provenance, coordinates, voucher codes, haplotype and GenBank accession numbers.

Specimen Voucher	Species name	Sample	Place of origin	Country	Haplotype	GenBank a.n.	Source
MIB:ZPL:07776	<i>Hystrix cristata</i>	Tissue	Prata, Grosseto, Tuscany	Italy	HAP1	LT746352	this study
MIB:ZPL:07777	<i>Hystrix cristata</i>	Quills	Monti Dauni, Foggia, Apulia	Italy	HAP1	LT746353	this study
MIB:ZPL:07778	<i>Hystrix cristata</i>	Quills	Monti Lepini, Rome, Latium	Italy	HAP1	LT746354	this study
MIB:ZPL:07779	<i>Hystrix cristata</i>	Quills	Veleggio Lomellina, Pavia, Lombardy	Italy	HAP1	LT746355	this study
MIB:ZPL:07780	<i>Hystrix cristata</i>	Quills	Saluzzo, Cuneo, Piedmont	Italy	HAP1	LT746356	this study
MIB:ZPL:07781	<i>Hystrix cristata</i>	Quills	Torrelvicino, Vicenza, Veneto	Italy	HAP1	LT746357	this study
MIB:ZPL:07782	<i>Hystrix indica</i>	Quills	Hula Valley	Israel	HAP2	LT746358	this study
MIB:ZPL:07783	<i>Hystrix indica</i>	Tissue	Adiyaman	Turkey	HAP3	LT746359	this study
WF-RFDF-11	<i>Hystrix indica</i>	–	–	India	HAP6	JN714177	GenBank
WF-RFDF-15-1	<i>Hystrix indica</i>	–	–	India	HAP5	JN714182	GenBank
WF-RFDF-15-2	<i>Hystrix indica</i>	–	–	India	HAP5	JN714183	GenBank
WF-RFDF-15-3	<i>Hystrix indica</i>	–	–	India	HAP5	JN714184	GenBank
MIB:ZPL:0784	<i>Hystrix</i> sp.	Quills	Abbadia di Fiastra, Macerata, Marche	Italy	HAP4	LT746360	this study

approaches based, for example, on High Resolution Melting through Real-Time PCR (Ramón-Laca et al., 2014).

The conservation status of the crested porcupine represents a European paradox. Despite most evidences suggest that it might have been introduced to Italy in Early mediaeval times (Bertolino et al., 2017; Trucchi et al., 2016), a few dissent voices suggesting a native origin are still present (see Bertolino et al. (2017) for a review). This species is listed in Berne Convention (all. II) and Habitat Directive (all. IV), as well as protected by the Italian law since 1978 (Italian National Law n. 968/1977). Starting from the 1970s, crested porcupines have undergone a remarkable range expansion in Italy (Mori et al., 2013). In the province of Macerata, where the *H. indica* sample was found, the population of porcupines started to become consistent and stable after 1976 (for the current distribution see Supplemental Figure S2). Our findings showed for the first time the presence of free-ranging Indian porcupine in this area of Central Italy. This species is quite common in Zoological Gardens and other Wildlife Centres (at least in 4 out of 23 structures surveyed in Italy). Hybridization between Asiatic and African porcupine species has been recorded in captivity (Mohr, 1965), so we cannot exclude that hybridization with locally escaped individuals might occur. In the Sinai peninsula, the opposite situation has occurred, with the presence of potentially introduced *H. cristata* overlapping with native, rare *H. indica* (Saleh and Basuony, 1998; Trucchi et al., 2016). The skull morphology of three individuals collected in this area (Saleh and Basuony, 1998), showed intermediate features between *H. cristata* and *H. indica*, therefore suggesting a possible hybridization which needs to be confirmed. A further step of monitoring would require the development of DNA-based tools, such as microsatellite or SNPs markers to properly distinguish possible hybrids from their parental species (Randi et al., 2001; Randi and Lucchini, 2002).

The management of the crested porcupine in Italy is controversial, as the species is widely poached for its meat (Lovari et al., 2016), and it is considered to be responsible for crop and riverbank damages (Bertolino et al., 2017; Laurenzi et al., 2016). This is somehow in contrast with the level of legal protection deserved to the species. Management strategies would necessarily change if a different porcupine species, surely introduced, would similarly affect human economy. We have no current evidence on the number of free-ranging *H. indica* individuals in the Macerata area. Immediate actions for the monitoring of occurrence distribution and breeding evidences of the species is necessary to control its potential expansion. Further attention by government and international agencies should be deserved to importation and detection of potentially dangerous animal species in zoological gardens and wildlife centres. We recommend to continue the monitoring of the range expansion of the crested porcupine in Italy and the collection of quills (especially where doubtful individuals are observed) for DNA analyses.

## References

Abiadh A., Chetoui M., Lamine-Cheniti T., Capanna E., Colangelo P. 2010. Molecular phylogenetics of the genus *Gerbillus* (Rodentia, Gerbillinae): implications for systematics, taxonomy and chromosomal evolution. *Mol. Phil. Evol.* 56: 513–518.

- Aulagnier S., Haffner P., Mitchell-Jones A.J., Moutou F., Zima J. 2010. Guide des Mammifères d'Europe, d'Afrique du Nord et du Moyen-Orient. Delachaux & Niestlé SA (Ed.), Paris, France: pp. 1–248. [in French]
- Barthelmeß E. 2006. *Hystrix africae australis*. *Mammal. Spec.* 788: 1–7.
- Bensasson D., Zhang D. X., Hartl D. L., Hewitt G. M. 2001. Mitochondrial pseudogenes: evolution's misplaced witnesses. *Trends Ecol. Evol.* 16: 314–321
- Bertolino S., Colangelo P., Mori E., Capizzi D. 2015. Good for management, not for conservation: an overview of research, conservation and management of Italian small mammals. *Hystrix* 6: 25–35. doi:10.4404/hystrix-26.1-10263
- Castiglia R., Aloise G., Amori G., Annesi F., Bertolino S., Capizzi D., Mori E., Colangelo P. 2016. The Italian peninsula hosts a divergent mtDNA lineage of the water vole, *Arvicola amphibius* s.l., including fossorial and aquatic ecotypes. *Hystrix* 27: 99–103. doi: 10.4404/hystrix-27.2-11588
- Colangelo P., Civitelli M.V., Capanna E. 2001. Morphology and chromosomes of *Tatera Lataste* 1882 (Rodentia Muridae Gerbillinae) in West Africa. *Tropical Zool.* 14: 243–253.
- Colangelo P., Leirs L., Castiglia R., D'aes M., Meheretu Y., Verheyen E. 2009. New data on the distribution and phylogenetic position of *Mastomys awashensis* (Rodentia, Muridae). *Mamm. Biol.* 75: 459–462.
- Corti M., Castiglia R., Colangelo P., Capanna E., Beolchini F., Bakele A., Ogue N.O., Makundi R.H., Sichilima A.M., Leirs H., Verheyen W., Verhagen R. 2005. Cytotaxonomy of rodent species from Ethiopia, Kenya, Tanzania and Zambia. *Belg. J. Zool.* 135: 197–216.
- Corbet G.B., Jones L.A. 1965. The specific characters of the crested porcupines, subgenus *Hystrix*. *Proc. Zool. Soc. Lond.* 144: 285–300.
- Ermakov O.A., Simonov E., Surin V.L., Titov S.V., Brandler O. V., Ivanova N. V., Borisenko A. V. 2015. Implications of hybridization, NUMTs, and overlooked diversity for DNA barcoding of Eurasian ground squirrels. *PLoS ONE* 10(1): e0117201. doi:10.1371/journal.pone.0117201
- Galimberti A., Spada M., Russo D., Mucedda M., Agnelli P., Crottini A., Ferri E., Martinoli A., Casiraghi M. 2012. Integrated operational taxonomic units (IOTUs) in echolocating bats: A bridge between molecular and traditional taxonomy. *Plos one* 7: e40122. doi: 10.1371/journal.pone.0040122
- Galimberti A., Sandionigi A., Bruno A., Bellati A., Casiraghi M. 2015. DNA barcoding in mammals: What's new and where next? *Hystrix* 26: 13–24. doi: 10.4404/hystrix-26.1-11347
- Gotzek D., Brady S.G., Kallal R.J., LaPolla J.S. 2012. The importance of using multiple approaches for identifying emerging invasive species: the case of the raspberry crazy ant in the United States. *PLoS ONE* 7: e45314. doi:10.1371/journal.pone.0045314
- Hall T.A. 1999. BioEdit: a user-friendly biological sequence alignment editor and analysis program for Windows 95/98/NT. *Nucleic acids symposium* 41:95–98.
- Hebert P.D.N., Cywinka A., Ball S.L., de Waard J.R. 2003. Biological identifications through DNA barcodes. *Proc. Royal Soc. Lond. B* 270: 313–321.
- Laurenzi A., Bodino N., Mori E. 2016. Much ado about nothing: assessing the impact of a problematic rodent on agriculture and native trees. *Mammal Res.* 61: 65–72.
- Lovari S., Corsini M.T., Guazzini B., Romeo G., Mori E. 2017. Suburban ecology of the crested porcupine in a heavily poached area: a global approach. *Eur. J. Wildl. Res.* doi: 10.1007/s10344-016-1060-7
- Mazzamuto M. V., Galimberti A., Cremonesi G., Pisanu B., Chapuis J-L., Stuyck J., Amori G., Su H., Aloise G., Preatoni D.G., Wauters L. A., Casiraghi M., Martinoli A., 2016. Preventing species invasion: A role for integrative taxonomy? *Integr. Zool.* 11: 214–228.
- Mohr E. 1965. *Allweltliche Stachelschweine*. Westarp Wissenschaften (Ed.), A. Ziemsen Verlag, Wittenburg Lutherstadt, Germany: pp. 1–163. [in German]
- Mori E., Sforzi A., Di Febbraro M. 2013. From the Apennines to the Alps: recent range expansion of the crested porcupine *Hystrix cristata* L., 1758 (Mammalia: Rodentia: Hystricidae) in Italy. *Ital. J. Zool.* 80:469–480.
- Padial J.M., Miralles A., de la Riva I., Vences M. 2010. The integrative future of taxonomy. *Front. Zool.* 7: 16.
- Pisanu B., Obolenskaya E.V., Baudry E., Lisovsky A.A., Chapuis J-L. 2013. Narrow phylogeographic origin of five introduced populations of the Siberian chipmunk *Tamias (Eutamias) sibiricus* (Laxmann, 1769) (Rodentia: Sciuridae) established in France. *Biol. Inv.* 15 1201–1207.
- Ramón-Laca A., Gleeson D., Yockney I., Perry M., Nugent G., Forsyth D.M. 2014. Reliable discrimination of 10 ungulate species using high resolution melting analysis of faecal DNA. *Plos one* 9: e92043. doi: 10.1371/journal.pone.0092043
- Randi E., Pierpaoli M., Beaumont M., Ragni B., Sforzi A. 2001. Genetic identification of wild and domestic cats *Felis silvestris* and their hybrids using Bayesian clustering methods. *Mol. Biol. Evol.* 18: 1679–1693.

- Randi E., Lucchini V. 2002. Detecting rare introgression of domestic dog genes into wild wolf (*Canis lupus*) populations by Bayesian admixture analyses of microsatellite variation. *Cons. Genet.* 3: 29–43.
- Saleh M.A., Basuony M. 1998. A contribution to the mammalogy of the Sinai Peninsula. *Mammalia* 62: 557–575.
- Skinner J., Smithers R.H.N. 1990. *Hystrix africaeaustralis*. In: The Mammals of the southern African subregion. University of Pretoria, Pretoria, South Africa: 93–97.
- Trucchi E., Gentile G., Sbordoni V. 2008. Development of primers to amplify mitochondrial DNA control region of Old World porcupines (subgenus *Hystrix*). *Mol. Ecol. Res.* 8: 1139–1141.
- Trucchi E., Sbordoni V. 2009. Unveiling an ancient biological invasion: molecular analysis of an old European alien, the crested porcupine (*Hystrix cristata*). *BMC Evol. Biol.* 9: 109.
- Trucchi E., Facon B., Gratton P., Mori E., Stenseth N.C., Jentoft S. 2016. Long live the alien: is high genetic diversity a pivotal aspect of crested porcupine (*Hystrix cristata*) long-lasting and successful invasion? *Mol. Ecol.* 25: 3527–3539.
- Walid M.F. 2011. The crested porcupine *Hystrix cristata* (Linnaeus, 1758) in Misurata, Libya. *Eur. J. Biol. Sci.* 3: 9–12.
- Wauters L.A., Amori G., Gippoliti S., Agnelli P., Galimberti A., Casiraghi M., Preatoni D.G., Martinoli A. 2017. New endemic mammal species for Europe: *Sciurus meridionalis* (Rodentia, Sciuridae). *Hystrix* 28(1): doi: 10.4404/hystrix-28.1-12015.
- Wiemers M., Fiedler K. 2007. Does the DNA barcoding gap exist? — A case study in blue butterflies (Lepidoptera: Lycaenidae). *Front. Zool.* 4:8.

Associate Editor: R. Caniglia

## Supplemental information

Additional Supplemental Information may be found in the online version of this article:

**Table S1** Analysis of the K2P distance matrix.

**Figure S2** Distribution of the crested porcupine in Central Italy.